



Patent
Attorney's Docket No. 021565-075



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

UTILITY PATENT
APPLICATION TRANSMITTAL LETTER

Box PATENT APPLICATION
Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Enclosed for filing is the utility patent application of Frank Meulewaeter, Marc Cornelissen, John Jacobs, Gerben Van Eldik and Michael Metzlaiff for METHODS AND MEANS FOR DELIVERING INHIBITORY RNA TO PLANTS AND APPLICATIONS THEREOF.

Also enclosed are:

- ☒ 1 sheet(s) of ☐ formal ☒ informal drawing(s);
- ☐ a claim for foreign priority under 35 U.S.C. §§ 119 and/or 365 is ☐ hereby made to filed in on ;
☐ in the declaration;
- ☐ a certified copy of the priority document;
- ☐ a General Authorization for Petitions for Extensions of Time and Payment of Fees;
- ☐ statement(s) claiming small entity status;
- ☐ an Assignment document;
- ☐ an Information Disclosure Statement; and
- ☒ Other: paper copy of Sequence Listing.
- ☒ An ☐ executed ☒ unexecuted declaration of the inventor(s)
☒ also is enclosed ☐ will follow.
- ☐ Please amend the specification by inserting before the first line the sentence --This application claims priority under 35 U.S.C. §§119 and/or 365 to filed in on ; the entire content of which is hereby incorporated by reference.--
- ☐ A bibliographic data entry sheet is enclosed.



21839

[X] The filing fee has been calculated as follows [X] and in accordance with the enclosed preliminary amendment:

C L A I M S					
	NO. OF CLAIMS		EXTRA CLAIMS	RATE	FEE
Basic Application Fee					\$690.00 (101)
Total Claims	52	MINUS 20 =	32	x \$18.00 (103)	\$576.00
Independent Claims	6	MINUS 3 =	3	x \$78.00 (102)	\$234.00
If multiple dependent claims are presented, add \$260.00 (104)					
Total Application Fee					\$1,500.00
If verified Statement claiming small entity status is enclosed, subtract 50% of Total Application Fee					
Add Assignment Recording Fee if Assignment document is enclosed					
TOTAL APPLICATION FEE DUE					\$1,500.00

- [] This application is being filed without a filing fee. Issuance of a Notice to File Missing Parts of Application is respectfully requested.
- [X] A check in the amount of \$ 1,500.00 is enclosed for the fee due.
- [] Charge \$ _____ to Deposit Account No. 02-4800 for the fee due.
- [X] The Commissioner is hereby authorized to charge any appropriate fees under 37 C.F.R. §§ 1.16, 1.17 and 1.21 that may be required by this paper, and to credit any overpayment, to Deposit Account No. 02-4800. This paper is submitted in duplicate.


Please address all correspondence concerning the present application to:

R. Danny Huntington, Esq.
Burns, Doane, Swecker & Mathis, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404.

Respectfully submitted,

BURNS, DOANE, SWECKER & MATHIS, L.L.P.

Date: April 18, 2000

By: 
Malcolm K. McGowan, Ph.D.
Registration No. 39,300

P.O. Box 1404
Alexandria, Virginia 22313-1404
(703) 836-6620

Methods and means for delivering inhibitory RNA to plants and applications thereof

Cross-Reference to Related Applications

- 5 This application is a continuation-in-part of U.S. Serial No. 09/294,022, filed April 20, 1999, the contents of which are incorporated herein by reference.

Field of the invention.

- 10 This invention relates to the field of functional genomics in plants, more particularly it relates to methods for the further identification and isolation of a nucleic acid with a nucleotide sequence of interest in a collection of preselected nucleic acid sequences correlated with a particular trait, preferably an agronomical important trait, using a kit of viral RNA vectors which allow systemic
- 15 spread of all components of the kit in a plant, wherein one of the viral RNA vectors comprises a library of gene-silencing constructs for the preselected nucleic acid sequences. The invention also relates to a method for modulating, preferably reducing, particularly eliminating the expression of a selected nucleic acid sequence, using the viral vector kit, whereby one of the vectors comprises
- 20 a gene-silencing construct for the selected nucleic acid sequence. The latter method may be used for validating the function of a nucleic acid sequence whose expression is correlated with the presence or absence of a specific trait in plants, but with otherwise unknown function. Preferably, one of the viral RNA vector components of the kit is a vector derived from a satellite virus.

25

Background art

- The recent, rapid expansion of available nucleic acid sequence information has
- 30 necessitated the development of methods for identifying the function of nucleic acid sequences, particularly transcribed nucleic acid sequences such as

expressed sequence tags, with unknown function, in an efficient and labor-cost effective way.

To identify the role of sequenced nucleic acids from plants of unknown function it is necessary to produce or identify plants in which those nucleic acids are either structurally or functionally inactivated. Plants wherein predetermined nucleic acid sequences are structurally inactivated can be generated using recombination technologies such as homologous recombination as described by Kempin et al. (1997) or using specific technologies such as the use of mixed duplex oligonucleotides (chimeraplasts) to generate specific mutations (as described in WO 96/22364 and WO 99/07865) . Alternatively, plants with a mutation in a predetermined nucleotide sequence can be identified by screening a saturated mutant library, such as but not limited to a T-DNA insertion library or a transposon insertion library (see e.g. Pereira and Aerts, 1998). These methodologies all require the generation of a large number of permanently altered plants, and thus are less amenable for application in high throughput methods. Moreover, the recovery of plants with recessive mutations in essential genes requires time-consuming breeding to maintain the plants in heterozygous state. Maintenance of dominant lethal mutations in essential genes is virtually impossible.

Plants with functionally inactivated predetermined nucleotide sequences can be generated in a straightforward way using methodologies wherein inhibitory RNA is generated, such as antisense or sense RNA.

The use of inhibitory RNA to reduce or abolish gene expression, also known as gene silencing, is well established in the art and is the subject of several reviews (e.g Baulcombe 1996, Stam et al. 1997, Depicker and Van Montagu, 1997). Several patent applications relate to the practical exploitation of gene silencing.

US 5,190,131 and EP 0 467 349 A1 describe methods and means to regulate or inhibit gene expression in a cell by incorporating into or associating with the

genetic material of the cell a non-native nucleic acid sequence which is transcribed to produce an mRNA which is complementary to and capable of binding to the mRNA produced by the genetic material of that cell.

- 5 EP 0 240 208 describes a method to regulate expression of genes encoded for in plant cell genomes, achieved by integration of a gene under the transcriptional control of a promoter which is functional in the host and in which the transcribed strand of DNA is complementary to the strand of DNA that is transcribed from the endogenous gene(s) one wishes to regulate.

10

EP 0 223 399 A1 describes methods to effect useful somatic changes in plants by causing the transcription in the plant cells of negative RNA strands which are substantially complementary to a target RNA strand. The target RNA strand can be a mRNA transcript created in gene expression, a viral RNA, or other RNA present in the plant cells. The negative RNA strand is complementary to at least a portion of the target RNA strand to inhibit its activity *in vivo*.

15

EP 0 647 715 A1 and US patents 5, 034,323, 5,231,020 and 5,283,184 describe methods and means for producing plants exhibiting desired phenotypic traits, by selecting transgenotes that comprise a DNA segment operably linked to a promoter, wherein transcription products of the segment are substantially homologous to corresponding transcripts of endogenous genes, particularly endogenous flavonoid biosynthetic pathway genes.

20

- 25 W0 93/23551 describes methods and means for the inhibition of two or more target genes, which comprise introducing into the plant a single control gene which has distinct DNA regions homologous to each of the target genes and a promoter operative in plants adapted to transcribe from such distinct regions RNA that inhibits expression of each of the target genes.

30

A major disadvantage of these technologies, which hampers the exploitation thereof in high throughput gene function discovery methods, is the intrinsic unpredictability and low occurrence of the gene silencing phenomenon.

5 Recently, Waterhouse et al. (1998) have described methods and means to make gene silencing in plants more efficient and predictable, by simultaneous expression of both sense and antisense constructs in cells of one plant. The sense and antisense nucleic acids may be in the same transcriptional unit, so that a single RNA transcript that has self-complementarity is generated upon
10 transcription.

Hamilton et al. (1998) describe improved silencing e.g. of tomato ACC-oxidase gene expression using a sense RNA containing two additional upstream inverted copies of its 5' untranslated region.

15 WO 98/53083 describes constructs and methods for enhancing the inhibition of a target gene within an organism, involving the insertion into the gene silencing vector of an inverted repeat of all or part of a polynucleotide region within the vector.

20 It should be clear however, that the use of inhibitory RNA as a tool in reversed genetics analysis of gene function via high throughput methods, whereby the inhibitory RNA is generated from gene-silencing constructs which are stably integrated in the genome of transgenic plants, suffers from the same drawbacks
25 as the methods wherein the nucleotide sequences are structurally inactivated.

EP 0 194 809 and US 5,500,360 suggest the use of viral RNA vectors to produce regulatory RNA such as anti-sense RNA.

30 Initial exploration of the use of viral vectors to deliver inhibitory RNA into cells of plants has been described by Chapman (1991). In this publication, gene silencing constructs comprising nucleotide sequences complementary to the

translated region of the GUS gene on a PVX derived viral vector were described. The experiments however, remained inconclusive as to whether gene silencing could be provoked using viral vectors for the production of inhibitory RNA.

- WO 93/03161 is directed toward recombinant plant viral nucleic acids and to hosts infected thereby. The non-native nucleic acid sequence which is transcribed may be transcribed as an RNA which is capable of regulating the expression of a phenotypic trait by an anti-sense mechanism.
- English et al., 1996 describe the suppression of the accumulation of a viral vector comprising a foreign nucleotide sequence in transgenic plants exhibiting silencing of nuclear genes comprising the same foreign nucleotide sequences, thus linking gene silencing and viral vectors, albeit in a reverse way as envisioned here.
- Kumagai et al. 1995 (PNAS 92, 1679-1683) described the inhibition of phytoene desaturase gene by viral delivery of antisense RNA.
- WO 95/34668 suggests the use of genetic constructs based on RNA viruses which replicate in the cytoplasm of cells to provide inhibitory RNA, either antisense or co-suppressor (sense) RNA.
- Baulcombe et al. (1998) and Ruiz et al. (1998) describe virus-induced gene silencing of the endogenous phytoene desaturase gene (PDS) or of a green fluorescent protein transgene (GFP) in plants, using potato virus X derived vectors carrying inserts homologous to PDS and GFP, respectively. The authors further suggested that virus-induced gene silencing may develop into a novel assay of gene function, by introducing a fragment of the genome of a viral vector and inferring the function of the gene from the symptoms of the infected plants exhibiting gene silencing.

The described methods for identification of the function of a gene with known nucleotide sequence however, have drawbacks and limitations. In the first place, the applicability of the mentioned viral RNA vector based gene silencing methods on larger scale is in practice limited to the identification of genes with essential functions or genes with macroscopically visible phenotypes. Secondly, all methods employ viral vectors which are capable of autonomous replication in plant cells and cell-to-cell movement, whereby care has to be taken not to inactivate the essential functions required for these functions. This may particularly be a disadvantage when tailoring these methods to the needs of particular plants, such as crop plants, by developing new viral vectors more apt for replication and systemic spread in the plants.

The prior art is thus deficient in the lack of efficient methods for large scale identification of the function of nucleic acids with known nucleotide sequence, or for the isolation of the genes of interest from a pool of genes with known nucleotide sequence, but unknown function.

Brief Description of the figures

Figure 1 is a schematic representation of the viral RNA vectors used in the Examples. MP: movement protein; CP: coat protein; OAS: origin of assembly. The open reading frames are indicated by boxes. Each original viral genome is characterized by a specific pattern.

▨: tobacco mosaic virus; ▩ tobacco necrosis virus; ▤ satellite tobacco mosaic virus; ■ satellite tobacco necrosis virus.

Summary of the invention

The invention provides a method for isolating genes involved in the determination of a trait or a phenotype of a plant species, comprising identifying a set of nucleic acids sequences of genes, whose expression is correlated with a trait of interest; creating a library of gene silencing constructs targeted or adapted to the nucleotide sequence of the identified nucleic acids in a viral RNA vector which is capable of replication inside plant cells and optionally, movement

between plant cells of a plant; infecting a collection of individual plants of the same plant species with the library of gene silencing constructs whereby each plant is infected with at least one member of the library; identifying a plant wherein the trait or phenotype is altered using an assay adapted to that trait or phenotype; and isolating the gene involved in the determination of the trait or phenotype in the plant species, from the library, based on the nucleotide sequence to which the gene silencing construct in the identified plant was targeted. Preferably the viral RNA vector is capable of autonomous replication inside plant cells and optionally autonomous movement between plant cells and particularly the viral RNA vector is derived from cowpea mosaic virus.

Alternatively, the viral RNA vector, is derived from a satellite RNA virus, preferably satellite tobacco mosaic virus or satellite tobacco necrosis virus, particularly it further comprises an origin of assembly of tobacco mosaic virus, and is capable of replication inside plant cells and optionally movement between plant cells when the required factors are supplemented *in trans*, preferably by infection with a helper virus, preferably tobacco mosaic virus or a helper virus which is derived from tobacco necrosis virus and comprises a gene encoding a coat protein gene of tobacco mosaic virus, and optionally, a movement protein of tobacco mosaic virus.

Gene-silencing constructs comprised within the viral RNA vector may comprise antisense RNA or sense RNA, preferably they may comprise an inverted repeat. Particularly the gene-silencing constructs comprise complementary stretch of at least 50, preferably at least 100 nucleotides of sense and antisense RNA. Especially preferred are gene-silencing constructs comprising at least two copies of part of the nucleotide sequences of the collection of nucleic acids, the copies being in inverted repeat.

It is another object of the invention to provide a method for the isolation or selection of a nucleic acid with a specific function from a collection of nucleic acids, wherein the collection of nucleic acids is characterized by the fact that

variation in the expression pattern of the nucleic acids is correlated with variation in a trait or phenotype of a plant harboring the nucleic acids comprising the steps of creating a library of gene silencing constructs targeted or adapted to the nucleotide sequence of the identified nucleic acids in a viral RNA vector which is capable of replication inside and optionally movement between plant cells; infecting a collection of plants with the library of gene silencing constructs, whereby each plant is infected with one member of the library; identifying plants with altered trait or phenotype using an assay adapted to the trait or phenotype under investigation; and optionally isolating the nucleic acid with the specific function from the identified plant with altered trait or phenotype.

It is yet another object of the invention to provide a method for determining the function encoded by a nucleic acid comprising a known nucleotide sequence in a plant, comprising the steps of providing a viral RNA vector derived from a satellite RNA virus, comprising a gene-silencing construct targeted to a gene comprising the known nucleotide sequence; infecting or inoculating the plant with the chimeric viral RNA vector and a corresponding helper virus or helper virus RNA; and identifying an altered trait or phenotype of the co-infected plant.

The invention further provides a method for isolating essential genes from a plant, comprising creating a library of random gene-silencing constructs, preferably by cloning random DNA fragments of the plant or by cloning random cDNA fragments, particularly duplicated cDNA fragments in inverted repeat in a cDNA copy of the viral RNA vector derived from a satellite RNA virus, preferably STMV or STNV, particularly a viral RNA vector comprising an origin of assembly of TMV; infecting a plant with individual members of the library and with a corresponding helper virus, preferably tobacco mosaic virus or a helper virus derived from tobacco necrosis virus and comprising the coat protein of tobacco mosaic virus and optionally the movement protein; identifying plants developing gene-silencing-construct-associated necrosis and optionally isolating the viral RNA vector from the necrotized tissue.

It is yet another object of the invention to provide a method for introduction of inhibitory RNA, preferably sense or antisense RNA, particularly inhibitory RNA comprising an inverted repeat, especially inhibitory RNA comprising a complementary stretch of at least 50, preferably at least 100 nucleotides of sense and antisense RNA, into plant cells, preferably into the cytoplasm of plant cells, comprising introducing into a plant cell, a viral RNA vector comprising the inhibitory RNA or comprising a chimeric nucleic acid which when transcribed yields the inhibitory RNA, wherein the viral RNA vector is derived from a satellite RNA virus, preferably STMV or STNV, particularly a STMV - derived or STNV derived RNA vector comprising an origin of assembly from tobacco mosaic virus; and introducing into the same plant cell, a corresponding helper virus, preferably tobacco mosaic virus or a chimeric helper virus derived from tobacco necrosis virus and comprising a coat protein gene of tobacco mosaic virus and optionally, a movement protein encoding gene of tobacco mosaic virus.

The invention further provides a kit for introduction of inhibitory RNA, preferably sense or antisense RNA, particularly inhibitory RNA comprising an inverted repeat, especially inhibitory RNA comprising a complementary stretch of at least 50, preferably at least 100 nucleotides of sense and antisense RNA in the cytoplasm of a plant cell comprising 1) a viral RNA vector derived from a satellite RNA virus, comprising the inhibitory RNA or which comprise a chimeric nucleic acid which when transcribed yields the inhibitory RNA; and 2) a corresponding helper virus.

A particularly preferred kit comprises 1) a viral RNA vector derived from a satellite tobacco mosaic virus, comprising an origin of assembly of tobacco mosaic virus, further comprising or encoding the inhibitory RNA; and 2) a corresponding helper virus derived from a tobacco mosaic virus.

Another particularly preferred kit comprises 1) a viral RNA vector derived from satellite tobacco necrosis virus, especially STNV-2 or STNV-C and comprising an origin of assembly of tobacco mosaic virus further comprising or encoding the

inhibitory RNA; and 2) a corresponding helper virus derived from tobacco necrosis virus, particularly TNV-A or TNV-D, which comprises a coat protein gene of tobacco mosaic virus and optionally a movement protein gene of tobacco mosaic virus.

5

Detailed description of preferred embodiments.

The following definitions apply throughout this application, unless otherwise specified.

10 As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or
15 amino acids than the actually cited ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA region which is functionally or structurally defined, may comprise additional DNA regions etc.

As used herein, "a trait of a plant" indicates a phenotype which is the combined
20 result of the coordinated expression of a number of genes. Typical traits include yield, heterosis, drought-resistance, stress-resistance, high or low temperature-resistance, vigor, seed yield, plant habitat, architecture etc. Typically, a trait of a plant is named after its intended appearance.

25 A used herein a "phenotype" of a plant refers to any quantitative or qualitative characteristic of that plant, be it morphological (including macroscopic and microscopic characteristics), biochemical (including the presence, absence or concentration of particular metabolites or molecules) functional or other.

30 The term "gene" means any DNA or RNA fragment comprising a region (the "transcribed region") which is transcribed into a RNA molecule (e.g., a mRNA) in a cell, operably linked to suitable regulatory regions, e.g., a plant-expressible

promoter. A gene may thus comprise several operably linked fragments such as a promoter, a 5' leader sequence, a coding region, and a 3' region comprising a polyadenylation site. A plant gene endogenous to a particular plant species or virus (endogenous plant or virus gene) is a gene which is naturally found in that plant species or virus, or which can be introduced in that plant species by breeding techniques such as conventional breeding techniques. A chimeric gene is any gene which is not normally found in a plant species or, alternatively, any gene in which the promoter is not associated in nature with part or all of the transcribed DNA region or with at least one other regulatory region of the gene.

The term "expression of a gene" refers to the process wherein a DNA or RNA region which is operably linked to appropriate regulatory regions, particularly to a promoter, is transcribed into an RNA which is biologically active i.e., which is either capable of interaction with another nucleic acid or which is capable of being translated into a biologically active polypeptide or protein. A gene is said to encode an RNA when the end product of the expression of the gene is biologically active RNA, such as e.g. an antisense RNA, a ribozyme or a replicative intermediate. A gene is said to encode a protein when the end product of the expression of the gene is a biologically active protein or polypeptide. In addition to the above defined elements, a gene may further comprise elements for cap-independent translation such as an internal ribosome entry sequence or the first and second translation enhancing elements as defined in WO 97/49814.

As used herein the terms "gene-silencing" or "inhibitory" are not to be interpreted as meaning a complete abolishing of the expression of the target gene(s) but also includes any reduction in expression, measured either as a reduction in transcription and/or translation, as a reduction in the accumulation of transcripts or translation products such as proteins, or as a reduction in the phenotypic expression of the target gene.

The term "reduction of phenotypic expression" refers to the comparison of the phenotypic expression of the nucleic acid of interest in the eukaryotic cell in the presence of the inhibitory RNA or gene-silencing constructs of the invention, to the phenotypic expression of the nucleic acid of interest in a similar eukaryotic cell in the absence of the inhibitory RNA or gene-silencing constructs of the invention. The phenotypic expression in the presence of the inhibitory RNA of the invention should thus be lower than the phenotypic expression in absence thereof, preferably be only about 25%, particularly only about 10%, more particularly only about 5% of the phenotypic expression in absence of the inhibitory RNA, especially the phenotypic expression should be completely inhibited for all practical purposes by the presence of the inhibitory RNA or the gene-silencing construct encoding such an RNA.

A reduction of phenotypic expression of a nucleic acid where the phenotype is a qualitative trait means that in the presence of the inhibitory RNA, the phenotypic trait switches to a different discrete state when compared to a situation in which such inhibitory RNA is absent. A reduction of phenotypic expression of a nucleic acid may thus also be measured as a reduction in transcription of (part of) that nucleic acid or reduction in the level of transcript, a reduction in translation of (part of) that nucleic acid or reduction in the level of translation products, or a reduction of the effect the presence of the transcribed RNA(s) or translated polypeptide(s) have on the eucaryotic cell or the organism, and will ultimately lead to altered phenotypes. It is clear that the reduction in phenotypic expression of a nucleic acid of interest, may be accompanied by or correlated to an increase in a phenotype or trait.

In one embodiment of the invention, a method is provided to identify and isolate genes involved in the determination of a trait or a phenotype of a plant. To this end, nucleic acid sequences are identified whose expression is correlated with the trait and/or phenotype of interest. Methods and means are available in the art for the almost simultaneous identification and/or isolation of a large number, if not the predominant part, of nucleotide sequences whose expression,

particularly whose transcription, is influenced subsequent to a stimulus corresponding to the trait to be investigated, in comparison with expression of these nucleotide sequences in a control plant. Such methods include but are not limited to differential display methods, such as the gel -based RNA differential display methods described by (Prahasar et al. 1996)

As a result of these methods, a collection of at least partially characterized nucleotide sequences with altered expression in response to a particular stimulus is identified. Typically, however, the application of such methods does not allow to discriminate between genes whose altered expression is directly caused by the stimulus, and those who are further downstream in the chain of events and are only indirectly influenced by the stimulus and a further selection amongst the obtained collection of nucleotide sequences will be required. Even less do these methods allow to predict whether the inverse relationship also holds, i.e. whether influencing the expression of genes with particular nucleotide sequences, identified in the above mentioned way, also influences the trait of interest. A further validation of the obtained sequences is thus required, and preferably one which immediately verifies the above mentioned inverse relationship. To achieve this goal in a efficient and cost-effective way, a library of gene-silencing constructs may be created in a viral RNA vector which is capable of replication inside plant cells and movement between cells of a plant, so as to assure an efficient systemic spread within a plant which has been infected with a clone of such a library. The created library of gene-silencing constructs comprised within a viral RNA vector is then used to infect a representative number of plants in such a way that at each plant is infected by at least one member (one clone) of the library. The infected plants can than be analyzed to identify those plants which exhibit alterations in the trait under investigation, using an assay which is adapted to the trait under investigation. It is clear that this fine-tuning of assay and trait under investigation may be an important advantage over the existing methods for high throughput analysis, particularly when analyzing traits and/or phenotypes which do not result in macroscopically visible alterations, such as but not limited to modifications in specific metabolic pathways or alterations which are only detectable under

specific conditions (e.g. heat, stress, drought-tolerance, pathogen-infection, application of specific herbicides or insecticides etc.).

The subset of nucleic acid sequences may also be identified on the basis of the presence of a particular signature characteristic of a class of proteins, such as but not limited to a kinase-specific domain, a binding motif etc.

The gene-silencing construct may then be isolated from the library or from the plant exhibiting the altered trait or phenotype, and be used to isolate the corresponding gene based on the nucleotide sequence towards which the gene silencing construct was targeted.

In a preferred embodiment, the viral RNA vector is capable of autonomous replication inside plant cells and autonomous movement between plant cells.

Such viral RNA vectors are known in the art and may be based on Potato Virus X as described e.g. in WO 93/03161, WO95/34668, Ruiz et al. (1998)

In a particularly preferred embodiment, the used viral RNA vector is derived from cowpea mosaic virus. Wellink et al. (1998) have described the use of a viral vector derived from this RNA virus for the expression of GFP in plants and demonstrated that the virus and RNA vectors derived thereof have an excellent capacity for spreading throughout an infected plant, particularly *Nicotiana benthamiana*. CPMV is an icosahedral virus with a bipartite RNA genome, consisting of a longer and a shorter RNA. Wellinck et al. have demonstrated that it is possible to incorporate extra genetic information in the shorter RNA, by inserting the GFP coding region in frame into the viral encoded polyprotein. For the purpose of the herein described methods, it is preferred that the inhibitory RNA encoding nucleic acid be inserted downstream of the polyprotein encoding open reading frame.

The inventors have obtained for the first time indications that gene-silencing may be obtained in plant cells such as protoplasts, using a viral vector derived from

a satellite virus comprising a β -1,3-glucanase coding region, in a co-infection experiment with a helper virus of the satellite virus

In another preferred embodiment, the viral RNA vector is capable of replication and cell-to-cell movement, only when the required functions are provided *in trans*. Particularly preferred, is the use of a satellite virus derived RNA vector, which can replicate in plant cells and spread throughout the plant, when a corresponding helper virus is present.

As used herein, "a satellite virus" indicates an RNA virus, preferably a single stranded RNA virus, the RNA genome of which is capable of replicating in a plant cell and being encapsidated by coat protein molecules to form a virus particle or virion, only when provided externally with any number of required essential functions therefor. By "externally provided" is meant that such functions are not encoded by the satellite viral genome. Satellite viruses thus depend upon external provision of essential functions, and may lack the capacity to encode functional replicase, movement protein, or other essential functions required to complete their life cycle inside a plant cell. In a natural situation, such essential functions are usually provided by an autonomously replicating virus or so-called helper virus .

Satellite viruses useful for the present invention may include wild type isolates, but also encompassed by this definition are variants which result in reduced or minimal symptoms when infected on a host plant, particularly when co-inoculated with a corresponding helper virus. The definition also includes synthetic satellite viruses such as defective viruses and chimeric satellite viruses.

A "viral RNA vector derived from a satellite virus" should at least include cis elements from a satellite virus which are recognized by an externally provided replicase, and an origin of assembly allowing encapsidation by the provided coat protein. Preferably, the viral RNA vector does not comprise a gene encoding a

functional coat protein, particularly it does not comprise the nucleotide sequence which is essentially similar to the nucleotide sequence encoding a coat protein gene. Particularly preferred are viral RNA vectors comprising an origin of assembly recognized by coat protein molecules from a rod-shaped virus, such as tobacco mosaic virus, since rod-shaped viruses do not exhibit the spatial constraints imposed on the size of genome by icosahedral viruses, thus allowing a larger number of additional nucleotides to be incorporated in the viral vector. Conveniently, the viral RNA vector comprises a number of unique or low-occurrence restriction recognition sites.

The use of viral RNA vectors derived from a satellite virus, additionally solves problems associated with the use of viral RNA vectors, such as reducing the size of the vectors, increasing versatility etc.

Particularly suited for the invention are viral RNA vectors derived from satellite tobacco mosaic virus comprising the origin of assembly (OAS) from TMV, preferably comprising the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5443 to the nucleotide at position 5518 or the nucleotide of SEQ ID No 5 from the nucleotide at position 5430 to the nucleotide at position 5505 (such as the nucleotide sequence of SEQ ID No 12) and wherein the coat protein encoding gene has been deleted. Also particularly suited for the invention are viral RNA vectors derived from satellite necrosis vector strain comprising the OAS from TMV and wherein most of the coat protein gene has been deleted. Non-limiting examples of viral RNA vectors, suitable for the invention are described hereinafter.

"A corresponding helper virus" as used herein, indicates those RNA viruses, preferably single stranded RNA viruses, which can supply the satellite virus or the derived viral RNA vector with the functions required *in trans* by that satellite virus or the derived viral RNA vector, to allow it to replicate in the cytoplasm of plant cells, and spread throughout an infected plant. Typically, corresponding helper viruses will provide the satellite virus or the vector derived thereof with a replicase (RNA dependent RNA polymerase) which recognizes the *cis*

sequences present on the satellite virus RNA, and will allow replication of the satellite virus genome or the derived vector. Other proteins which may typically be provided by the helper virus are movement proteins, allowing inter alia, the plasmodesmata-mediated spread of viral particles from cell to cell. For satellite

5 viruses or viral RNA vectors derived thereof which lack a functional coat protein encoding gene, corresponding helper viruses may also provide a functional coat protein. Preferably, the corresponding helper virus will be capable of autonomous systemic spread in an infected plant. However, such a systemic spread seems not to be a prerequisite for efficient gene silencing. Functions

10 required *in trans* for one particular viral RNA vector may be supplied *in trans* by different corresponding helper viruses.

It is clear that the corresponding helper viruses may be wild type isolates of RNA viruses, preferably single-stranded RNA viruses such as the tobamoviruses or

15 necroviruses. Particularly preferred are rod-shaped RNA viruses such as tobamoviruses including tobacco mosaic virus and the related tobamoviruses such as ribgrass mosaic virus, turnip vein clearing virus, chine rape mosaic virus, oilseed rape mosaic virus.

20 When TMV can be used as a helper virus, it can also be replaced by one of the closely related tobamoviruses mentioned above, particularly when using the viral vectors in particular plant species.

Also encompassed by the methods and means of the invention are variants of

25 such wild type isolates, preferably variants or mutants which develop minimal symptoms when inoculated on host plants or when co-infected with a corresponding satellite virus or RNA vector derived thereof. Further preferred helper viruses may be variants or mutants of wild type isolates which have an extended host range such as tobamoviruses which can replicate and spread in

30 corn or brassicae.

However, corresponding helper viruses may also be chimeric or hybrid viruses, wherein part of the viral genome has been replaced by a foreign nucleic acid, particularly wherein part of the viral genome has been replaced by a nucleic acid derived from another viral genome, preferably a part comprising a nucleotide sequence encoding a movement protein, or a part comprising a nucleotide sequence encoding a coat protein. E.g. when using a necrovirus such as TNV, it may be advantageous to insert a movement protein encoding region, preferably a movement protein derived from a tobamovirus such as TMV, particularly the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 4903 to the nucleotide at position 5709, so as to ensure spreading of the viral particles beyond the infected leaf. However, spreading of the helper virus or the viral RNA vector is not essential for efficient inactivation of expression of the target genes throughout the plant, as was found by the inventors. Also when using e.g. a necrovirus such as TNV, it may be further advantageous to replace the coat protein coding region of the necrovirus by a coat protein coding region of a rod-shaped virus, such as TMV, particularly the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5712 to the nucleotide at position 6191. It goes without saying that an appropriate origin of assembly for the substituted coat protein has to be incorporated in the genome of the chimeric helper virus.

In the above described example however, the OAS of TMV is conveniently located within the movement protein coding region. Non-limiting examples of corresponding helper viruses will be described hereinafter.

It should be clear that whenever it is stated that plants are co-infected or infected with a viral RNA vector and a corresponding helper virus, it is equal whether the helper virus is inoculated before, after or simultaneous with the viral RNA vector, provided however that there is a reasonable time limit between infection of the viral RNA vector or the corresponding helper virus.

Alternatively, the required functions *in trans* for the replication and movement of the viral RNA vector may be provided from the expression of chimeric genes, encoding a replicase (RNA dependent RNA polymerase) and/or a movement

protein and/or a functional coat protein, integrated in the genome of the test plants.

Preferred kits to deliver inhibitory RNA or gene-silencing constructs to plant cells to be used in the herein disclosed methods comprise a viral RNA vector derived from a satellite RNA virus, particularly from satellite tobacco necrosis vector (STNV) or satellite tobacco mosaic virus (STMV) and a corresponding helper virus, particularly a rod-shaped corresponding helper virus, wherein the viral RNA vector comprises a gene-silencing construct.

In a preferred embodiment the kit comprises a viral RNA vector derived from satellite tobacco necrosis vector, preferably comprising the cis-elements required for replication, particularly comprising the nucleotide sequence of SEQ ID No 3 from the nucleotide at position 1 to the nucleotide at position 32 and the nucleotide sequence of SEQ ID No 3 from the nucleotide at position 738 to the nucleotide at position 1245, wherein an origin of assembly of tobacco mosaic virus has been inserted, preferably comprising the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5443 to the nucleotide at position 5518 or comprising the nucleotide sequence of SEQ ID No 5 from the nucleotide at position 5430 to the nucleotide at position 5505, or comprising the nucleotide sequence of SEQ ID No 12 and wherein said helper virus is derived from tobacco necrosis virus, preferably with a nucleotide sequence of SEQ ID No 1, and comprises a gene encoding the movement protein of tobacco mosaic virus, preferably with the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 4903 to the nucleotide at position 5709 or with the nucleotide sequence of SEQ ID No 15 from the nucleotide at position 479 to the nucleotide at position 1285 and a gene encoding the coat protein of tobacco mosaic virus, preferably with the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5712 to the nucleotide at position 6191 or with the nucleotide sequence of SEQ ID No 15 from the nucleotide at position 1288 to the nucleotide at position 1767.

Preferred combinations are those kits wherein the viral RNA vector is derived from STNV-1 or STNV-2 strains (as disclosed by Ysebaert et al. 1980; Genbank Accession number M10388 or Danthinne et al., 1991 Genbank Accession M64479) and the helper virus is TNV-A (Meulewaeter et al 1990, SEQ ID No 1).

5 Other preferred combinations are those kits wherein the viral RNA vector is derived from STNV-C (Bringloe et al. (1998); Genbank Accession Nr AJ000898) and the corresponding helper virus is TNV-D (Coutts et al. (1991); Genbank Accession Nr D00942).

10 In another particularly preferred embodiment the kit comprises a viral RNA vector derived from satellite tobacco mosaic virus, preferably comprising the cis-elements required for replication, particularly comprising the nucleotide sequence of SEQ ID No 4 from the nucleotide at position 1 to the nucleotide at position 197 and the nucleotide sequence of SEQ ID No 4 from the nucleotide at position 604 to the nucleotide at position 1058 or comprising the nucleotide sequence of SEQ ID No 13 and the nucleotide sequence of SEQ ID No 14; and further comprising an origin of assembly of tobacco mosaic virus, preferably comprising the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5443 to the nucleotide at position 5518 or comprising the nucleotide sequence of SEQ ID No 5 from the nucleotide at position 5430 to the nucleotide at position 5505 or comprising the nucleotide sequence from SEQ ID No 12, and wherein said corresponding helper virus is a tobacco mosaic virus, particularly TMV-U1 (SEQ ID No 2) or TMV-U2 (SEQ ID No 5).

25 It will be clear to the person skilled in the art that viral RNA vectors may be generated conveniently by *in vitro* transcription methods from cDNA copies of the viral RNA. Likewise, infectious viral RNA for the corresponding helper viruses may be generated from cDNA copies of their genome. Libraries, viral vectors and corresponding helper viruses may also be maintained by replication in plant cells.

30

Methods to infect or inoculate plants and plant cells with viral RNA vectors, helper viruses and libraries comprised within viral RNA vectors are well within in the realm of the person skilled in the art and may be performed according to the methods described in Walkey (1985).

5

In one embodiment of the methods of the invention, plants are inoculated, e.g. with a solution containing the libraries of gene-silencing constructs in a viral vector, or with a solution containing a mixture of gene-silencing constructs in a viral vector and corresponding helper virus. The solution may further contain
10 additional compounds to improve inoculation and infection of the plants, such as, but not limited to abrasives, adherents, tensio-active products and the like.

15

Plants may be infected during different developmental stages, in order to maximize the phenotype under investigation. Also different parts of plants may be inoculated to optimize observation of the expected phenotype.

20

Although not intending to limit the scope of the invention to a particular mode of action, it is thought that the inhibitory RNA comprised within the viral RNA vector can exercise its inhibiting effect, provided there is a balance between RNA encapsidated in a virion and free RNA. It is thought that the balance between encapsidated and free RNA may be influenced by varying the sequence and position of an origin of assembly within the viral RNA vector.

25

However, the gene-silencing effect may be amplified by placing the inhibitory RNA encoding nucleic acid under control of a viral promoter, preferably a coat protein promoter, or a subgenomic promoter so that during the life cycle of the virus additional inhibitory RNA is generated or transcribed.

30

"Gene-silencing constructs" as used herein is to be interpreted as a nucleic acid, which when transcribed yield "inhibitory RNA" comprising or consisting of sense RNA or antisense RNA, or a combination of both comprising a nucleotide sequence which has at least 75%, preferably at least 80%, particularly at least 85%, more particularly at least 90%, especially at least 95% sequence identity

with or is identical to the nucleotide sequence whose expression is to be suppressed, or its complement. Further, the nucleotide sequence of the sense or antisense region should preferably be at least about 100 nucleotides in length, more preferably at least about 250 nucleotides, particularly at least about 500 nucleotides but may extend to the full length of the coding region of the gene whose expression is to be reduced.

For practical purposes in the application of the methods for high throughput screening or validation, the gene-silencing construct may be identical in sequence and length to the target nucleic acids, or they may be exactly complementary in sequence and identical in length to the target nucleic acids.

For the purpose of this invention the "sequence identity" of two related nucleotide or amino acid sequences, expressed as a percentage, refers to the number of positions in the two optimally aligned sequences which have identical residues (x100) divided by the number of positions compared. A gap, i.e. a position in an alignment where a residue is present in one sequence but not in the other is regarded as a position with non-identical residues. The alignment of the two sequences is performed by the Wilbur and Lipmann algorithm (Wilbur and Lipmann, 1983) using a window-size of 20 nucleotides or amino acids, a word length of 2 amino acids, and a gap penalty of 4. Computer-assisted analysis and interpretation of sequence data, including sequence alignment as described above, can be conveniently performed using commercially available software packages such as the programs of the Intelligenetics™ Suite (Intelligenetics Inc., CA) or the GCG Wisconsin Package.

It is clear for the person skilled in the art that the gene-silencing constructs may comprise at the same time sense and anti-sense RNA targeted towards the same nucleotide sequence whose expression is to be reduced. Preferably, the sense and antisense RNA are at least partly complementary to each other and capable of forming a stem-loop structure, since such a configuration has been shown to increase the efficiency of gene-silencing, both in occurrence and level

of gene-silencing (Waterhouse et al. 1998). In the most straightforward embodiment, at least part of the target nucleic acid, preferably the complete target nucleic acid, is cloned in duplicated form, whereby the two copies are in inverted repeat, preferably separated by an unrelated spacer nucleotide sequence.

The invention also aims at providing the herein described kits in their different embodiments. It is also an object of the invention to provide the kits comprising the helper viruses and viral RNA vectors described without the gene-silencing constructs or inhibitory RNA as well as their cDNA copies, whereby the cDNA copies are under control of a promoter (which can be used in *in vitro* transcription methods available in the art) such as but not limited to the promoters recognized by single subunit bacteriophage polymerase promoters (T7, T3, SP6 RNA polymerase specific promoters and the like).

The invention further relates to a method for identifying genes which are essential in plants comprising the following steps:

- a) A library of random gene-silencing constructs specific for the plant is created using a viral RNA vector which is derived from a satellite RNA virus, as herein defined including all its preferred embodiments. Preferably, the library is created in a cDNA copy of the viral vector and may be generated by inserting random DNA sequences, preferably at least about 100 nucleotides in length, particularly at least about 500 nucleotides in length. The random DNA sequences may be obtained from total DNA of a plant or may represent a subset of the genome of a plant, such as DNA derived from organelles (plastids, chloroplasts, mitochondria etc.) Alternatively, the library may be created by inserting cDNAs generated by reverse transcriptase from RNA, preferably mRNA obtained from said plant. The library may be normalized, e.g. as described in Takayuki et al. (1995). The library should preferably be large enough in size, i.e. contain a sufficient number of independent clones to cover the genome of the plant, according to the standards known in the art. In a preferred embodiment, the library may contain duplication of the

- inserted nucleic acid whereby the copies are in inverted repeat. The inserted nucleic acid may be cloned downstream of a viral promoter such as, but not limited to a coat protein gene promoter or a subgenomic promoter. It will further be clear to the person skilled in the art that the relative orientation of the inserted nucleic acid, in relation to the RNA vector is only of limited importance since either sense or antisense inhibitory RNA will be generated.
- 5 the inserted nucleic acid, in relation to the RNA vector is only of limited importance since either sense or antisense inhibitory RNA will be generated.
- b) Assay plants are infected with individual members of the library and also with a corresponding helper virus. Infection may proceed according to any of the methods mentioned herein. Clearly, the DNA copy of the library should be converted into an RNA copy according to any of the methods described herein, preferably prior to the infection of the assay plants.
- 10 converted into an RNA copy according to any of the methods described herein, preferably prior to the infection of the assay plants.
- c) Plants developing a gene-silencing-construct-associated phenotype are identified. As used herein, a "gene-silencing-construct-associated phenotype" is meant to indicate a phenotype which is not observed when performing a mock inoculation with a viral RNA vector without gene-silencing construct, in combination with a corresponding helper virus on a similar plant. Preferred phenotypes comprise chlorosis, necrosis or any phenotype, preferably a morphological phenotype indicating that the infected tissue is inhibited or dying or deteriorating.
- 15 mock inoculation with a viral RNA vector without gene-silencing construct, in combination with a corresponding helper virus on a similar plant. Preferred phenotypes comprise chlorosis, necrosis or any phenotype, preferably a morphological phenotype indicating that the infected tissue is inhibited or dying or deteriorating.
- 20d) Optionally, isolating the viral RNA vector from the tissue exhibiting the gene-silencing-construct-associated phenotype according to methods available in the art for isolation of virus. Preferably, the isolated viral RNA vector comprising the gene-silencing construct of interest should be re-assayed on fresh plants to confirm the observed phenotype. The viral RNA vector can of course also be recovered from the library if the infection of the plants was performed in an identity-preserving way.
- 25 course also be recovered from the library if the infection of the plants was performed in an identity-preserving way.
- e) The gene silencing construct or the nucleotide sequence information thereof, may then be used to recover the corresponding genomic or cDNA clone using methods available in the art (hybridization, PCR etc.)

As defined herein "essential genes" of a plant, are those genes which are necessary during the normal development of a plant. As defined, essential

genes may be essential for normal development only in particular developmental stages, or only in particular tissues or organs, such as e.g. flowers. Typically, inhibition of the expression of essential genes may have a lethal effect on a plant or part of a plant. Preferred essential genes are those genes which result in
 5 retardation or dying of seedlings when inhibited.

It will be clear for the person skilled in the art that if the inhibition of the target nucleic acid results in a dominant effect, as is the case for inhibition of the expression of essential genes, the described methods may be performed using
 10 infection of more than one viral RNA vector comprising a gene-silencing construct per plant. Care has to be taken to not dilute the phenotypic effect too much by infecting a too large number of different viral RNA vectors comprising differing gene-silencing constructs on the same plant. It is thought that optimally any number between one and five different inhibitory RNAs may be introduced
 15 in one plant cell.

In yet another embodiment of the invention, a method is provided for determining the function encoded by a nucleic acid comprising a known nucleotide sequence. This nucleotide sequence may have been obtained e.g. from a
 20 genome sequencing program, including expressed sequence tags sequencing programs. In order to unravel the function of that sequence, a gene-silencing construct or inhibitory RNA targeted towards said nucleotide sequence, as described in all its embodiments herein, may be introduced into a viral RNA vector derived from a satellite virus, as described herein, and used to inoculate
 25 a plant or being introduced into a plant cell, particularly into a protoplast, together with a corresponding helper virus, as described herein.

A large number of the embodiments described herein thus relate to a method for the introduction of inhibitory RNA in plant cells, comprising the steps of :

- 30a.) introducing into a plant cell, a viral RNA vector comprising inhibitory RNA or comprising a chimeric nucleic acid which when transcribed

yields the inhibitory RNA, wherein the viral RNA vector is derived from a satellite RNA virus; and

b.) introducing into the same plant cell, a corresponding helper virus.

- 5 The methods of the invention can be applied to essentially all plants for which viral vector and/or corresponding helper viruses are available. The methods of the invention are thought to be particularly suited for *Nicotiana* spp, particularly *N. tabacum*, *N. sylvestris*, *N. benthamiana*, and other *Solanaceae*, rice (*Oryza sativa*) corn (*Zea mays*), *Brassica* spp. , cotton (*Gossypium hirsutum*), wheat,
- 10 *Arabidopsis* spp., *Petunia* spp.

Also envisioned by the present invention are methods for developing an agronomically useful product, such as a herbicide or a transgenic plant using the herein described methods and means, further comprising the steps of inserting

15 a nucleic acid, involved in the determination of a particular plant trait, isolated by the methods of the invention, preferably under control of a foreign plant-expressible promoter, particularly under control of a controllable plant-expressible promoter into the genome of a plant , particularly a crop plant. When essential genes have been identified according to the methods described herein,

20 these essential genes or their encoded gene products, particularly the encoded proteins may be used in *in vitro* assays to identify compounds inhibiting the activity, particularly the enzymatic activity, which may be used as herbicides. Alternatively, a viral RNA vector encoding gene-silencing constructs targeted towards essential genes may be used as herbicidal compounds.

25

The following non-limiting Examples describe the construction of viral RNA vectors derived from satellite viruses, and uses thereof. Unless stated otherwise in the Examples, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, NY

30 and in Volumes 1 and 2 of Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, *Current Protocols*, USA. Standard materials and methods for plant

molecular work are described in *Plant Molecular Biology Labfax* (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

- 5 Throughout the description and Examples, reference is made to the following sequences:

SEQ ID No 1: nucleotide sequence of the genome of TNV-A

SEQ ID No 2: nucleotide sequence of the genome of TMV-U1 (Genbank
10 Accession Nr V01408).

SEQ ID No 3: nucleotide sequence of the genome of STNV-2

SEQ ID No 4: nucleotide sequence of the genome of STMV (Genbank
accession Nr. M25782).

SEQ ID No 5: nucleotide sequence of the genome of TMV-U2 (Genbank
15 Accession Nr M34077).

SEQ ID No 6: nucleotide sequence of the tomato phytoene desaturase (*pds*)
encoding cDNA (Genbank Accession Nr. X59948).

SEQ ID No 7: nucleotide sequence of the tobacco nitrate reductase (*nia-2*)
encoding cDNA (Genbank Accession Nr. X14059).

20 SEQ ID No 8: nucleotide sequence of the tobacco nitrite reductase (*nir-1*)
encoding cDNA (Genbank Accession Nr. X66145).

SEQ ID No 9: nucleotide sequence of the β -1,3-glucanase(*gn-1*) encoding
cDNA of *Nicotiana plumbagenifolia*.

SEQ ID No 10: nucleotide sequence of a green fluorescent protein (*gfp*)
25 encoding region.

SEQ ID No 11: nucleotide sequence of a β -glucuronidase (*gus*) encoding region.

SEQ ID No 12: nucleotide sequence of an origin of assembly of a TMV-U2
strain.

SEQ ID No 13: nucleotide sequence of the leader sequence of a STMV strain

30 SEQ ID No 14: nucleotide sequence of the trailer sequence of a STMV strain.

SEQ ID No 15: nucleotide sequence of part of the genome of a TMV-U2 strain
comprising movement protein and coat protein genes.

Examples

Example I : construction of the viral RNA vector kits

5 A plasmid vector for the synthesis of an infective hybrid TMV/TNV helper virus RNA is made using the following operationally linked elements:

- A T7 RNA polymerase promoter
- A nucleic acid comprising the nucleotide sequence from the nucleotides 1 to 2234 of TNV-A (nt 1 to 2234 of SEQ ID No 1), wherein the AUG codon at
10 nucleotides 2218-2220 mutated to a different codon
- A nucleic acid comprising the nucleotide sequence encoding the open reading frame for the movement protein of TMV-U1 (nt 4903-5709 of Genbank Accession Number V01408 or nt 4903-5709 of SEQ ID No 2 or nt 479-1285 of SEQ ID No 12)
- 15 • A nucleic acid comprising the nucleotide sequence from the nucleotides 2235 to 2612 of TNV-A (SEQ ID No 1)
 - A nucleic acid comprising the nucleotide sequence encoding the open reading frame for the coat protein of TMV-U1 (nt 5712-6191 of Genbank Accession Number V01408 or nt 5712-6191 of SEQ ID No 2 or nt 1288-
20 1767 of SEQ ID No 15)
 - A sequence comprising nucleotide 3444 to 3684 of TNV-A (nt 3444 to 3684 of SEQ ID No 1)

A plasmid vector for the synthesis of an infective hybrid TMV/STNV viral vector

25 RNA is made using the following operationally linked elements:

- A T7 RNA polymerase promoter
- A nucleic acid comprising the nucleotide sequence from nucleotide 1 to 32 of STNV-2 (nt 1 to 32 of SEQ ID No 3)
- A nucleic acid comprising the origin of assembly (OAS) of TMV-U1 (nt.
30 5443-5518 of Genbank Accession Number V01408 or nt 5443-5518 of SEQ ID No 2 or nt 1018 to 1094 of SEQ ID No 15)

- A nucleic acid comprising the nucleotide sequence from nucleotide 738 to 1245 of STNV-2 (nt 738 to 1245 of SEQ ID No 3)

A plasmid vector for the synthesis of an infective hybrid TMV/STMV viral vector

5 RNA is made using the following operationally linked elements:

- A T7 RNA polymerase promoter
- A nucleic acid comprising the nucleotide sequence from nucleotide 1 to 197 of STMV (Genbank Accession Number M25782; nt 1 to 197 of SEQ ID No 4) or the nucleotide sequence of SEQ ID No 13.
- 10 • A nucleic acid comprising the OAS of TMV-U1 (nt. 5443-5518 of Genbank Accession Number V01408; nt 5443 to 5518 of SEQ ID No 2; nt 1019-1094 of SEQ ID No 15) or of TMV-U2 (nt 5430-5505 of Genbank Accession Number M34077; nt 5430-5505 of SEQ ID No 5) such as the nucleotide sequence of SEQ ID No 12.
- 15 • A nucleic acid comprising the nucleotide sequence from nucleotide 604 to 1058 of STMV (Genbank Accession Number M25782; nt 604 to 1058 of SEQ ID No 4) or comprising the nucleotide sequence of SEQ ID No 14.

20 **Example 2. Feasibility demonstration using known endogenes or transgenes.**

To demonstrate the feasibility of the use of the viral kits described sub example 1 for functional knockout of specific endo- or transgenes in *Nicotiana* plants, one of the following DNA fragments is inserted in the TMV/STNV or TMV/STMV hybrid vector, immediately upstream or downstream of the TMV OAS:

- a fragment of the tomato pythoene desaturase (*pds*) cDNA (comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948)
- a fragment of the tobacco nitrate reductase (*nia-2*) cDNA (comprising 30 nucleotides 1103 to 2114 or nucleotides 5169 to 6497 of SEQ ID No 7 or Genbank Accession Number X14059)

- a fragment of the tobacco nitrite reductase (*nir-1*) cDNA (comprising nucleotide 650-1212 of SEQ ID No 8 or Genbank Accession Number X66145)
- a fragment of the β -1,3-glucanase (*gn-1*) cDNA of *Nicotiana plumbaginifolia* (SEQ ID No 9 or Genbank Accession Number X07280)
- a fragment comprising a nucleotide sequence from a green fluorescent protein (*gfp*) coding region (SEQ ID No 10)
- a fragment comprising a nucleotide sequence from nucleotide 1 to 600 of the β -glucuronidase (*gus*) coding region (SEQ ID No 11)

10

Infective chimeric transcripts are synthesized *in vitro*, using T7 RNA polymerase with the linearized plasmid DNAs of the described vectors as templates.

15

The TMV/STNV RNAs are mechanically inoculated on leaves of *Nicotiana benthamiana* or *Nicotiana tabacum* plants together with the TMV/TNV RNA, whereas the TMV/STMV RNAs are inoculated together with TMV-U2 virus particles or viral RNA.

The infected plants are scored for phenotypes, virus accumulation, and suppression of the homologous plant gene between 1 and 4 weeks after inoculation.

20

Plants infected with vectors containing the *pds* cDNA show a bleaching phenotype on infected leaves and silencing of the endogenous *pds* transcript. Plants infected with vectors containing the *nia-2* or *nir-1* cDNA show a chlorotic phenotype on infected leaves and silencing of the endogenous *nia-2* or *nir-1* transcript, respectively.

25

Plants infected with vectors containing the *gn-1* cDNA show silencing of the endogenous basic -1,3-glucanase transcript.

Upon infection with vectors containing the *gfp* sequence, transgenic plants that normally express a *gfp* transgene show silencing of the *gfp* transgene transcript and suppression of GFP fluorescence.

30

Upon infection with vectors containing the *gus* sequence, transgenic plants that normally express a *gus* transgene show silencing of the *gus* transgene transcript and suppression of GUS activity.

Example 3: Inactivation of phytoene desaturase in *Nicotiana benthamiana* using a TNV/STNV hybrid vector system.

A TNV/STNV hybrid vector system was used for the functional inactivation of a constitutively expressed endogenous plant gene . Therefore, the following STNV hybrid vectors have been constructed:

pIF9 carrying the following operationally linked elements:

- a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811;
- a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 32 of SEQ ID No 3 (STNV-2 leader);
- a nucleic acid comprising the origin of assembly (OAS) of TMV-U1 from nucleotide 5443 to 5518 of SEQ ID No 2 or Genbank Accession Number V01408;
- a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948;
- a nucleic acid comprising the nucleotide sequence from nucleotide 806 to 1418 of SEQ ID No 3 (STNV-2 trailer).

pIF12 carrying the following operationally linked elements:

- a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811;
- a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 32 of SEQ ID No 3 (STNV-2 leader);
- a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948;

- a nucleic acid comprising the nucleotide sequence from nucleotide 742 to 1354 of SEQ ID No 3 (STNV-2 trailer).

Infective chimeric transcripts have been synthesized in vitro using T7 RNA polymerase with the linearized plasmid DNAs of the described pIF9 and pIF12 hybrid vectors as templates using standard procedures. Control in vitro transcripts have been synthesized on linearized plasmid DNAs of the precursor plasmids of pIF9 and pIF12 without the *pds* fragment inserts and on linearized plasmid DNA of an infective clone of the STNV wild type and of a hybrid STNV vector carrying an insert of the *cat* gene.

The in-vitro transcripts have been mechanically inoculated onto leaves of four weeks old *Nicotiana benthamiana* plants together with the TNV helper virus.

All infected plants were continuously scored for *pds* inactivation, which resulted in a phenotype showing leaf bleaching. For all inoculations necrotic lesions were observed for the inoculated leaves after 2 days post inoculation (p.i.). Within a week, necrotic lesions also occurred in upper leaves indicating the systemic spread of the viruses in *N.benthamiana*. In most infected plants, the virus symptoms have been severe but plants survived for many weeks.

Only plants, which have been infected with the hybrid vectors pIF9 and pIF12 carrying *pds* fragments, showed on top of the virus symptoms additional phenotypic changes. Approximately 4 weeks p.i., green upper leaves, which did not show any virus symptoms, developed bleached spots scattered all over the leaves. The bleaching was progressive and was not accompanied by necrotic lesions. Within another three weeks, the size of the spots increased constantly and the color changed from pale green to yellow to pale white-yellow. These symptoms have never been observed in plants, on which TNV/STNV wild type or TNV/STNV-deletion mutant control inoculations were carried out. Thus, this phenotype indicates the functional knockout of *pds* in *N. benthamiana* plants after infection with pIF9 and pIF12 hybrid viral vectors.

In RNA gel blot analyses performed with total RNA preparations of green and bleached spots of upper leaves of plants showing a FKO phenotype no chimeric STNV virus RNA and only very low levels of TNV helper virus RNA could be detected. This indicates that virus induced gene silencing of pds in a specific tissue does not need to be accompanied by high levels of virus RNA.

Example 4: Inactivation of phytoene desaturase in *Petunia hybrida* using a TMV/STMV hybrid vector system.

A TMV/STMV hybrid vector system was used for the functional inactivation of a constitutively expressed endogenous plant gene. Therefore, the following STMV hybrid vectors have been constructed:

pVE293 carrying the following operationally linked elements:

- 15• a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811
 - a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 197 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV leader)
- 20• a nucleic acid comprising the origin of assembly (OAS) of TMV-U2 from nucleotide 5430 to 5505 of SEQ ID No 5 or Genbank Accession Number M34077 ("short" TMV OAS)
 - a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number
 - 25 X59948
 - a nucleic acid comprising the nucleotide sequence from nucleotide 604 to 1058 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV trailer)
 - a SP6 RNA polymerase promoter comprising nucleotide 143 to 124 of
 - 30 Genbank Accession Number X65308

pVE294 carrying the following operationally linked elements:

- a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811
- a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 197 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV leader)
- a nucleic acid comprising the origin of assembly (OAS) of TMV-U2 from nucleotide 5441 to 5849 of SEQ ID No 2 or Genbank Accession Number M34077 ("long" TMV OAS)
- 10 • a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948
- a nucleic acid comprising the nucleotide sequence from nucleotide 604 to 1058 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV trailer)
- 15 • a SP6 RNA polymerase promoter comprising nucleotide 143 to 124 of Genbank Accession Number X65308

Infective chimeric transcripts have been synthesized in vitro using T7 RNA polymerase with the linearized plasmid DNAs of the described pVE293 and pVE294 hybrid vectors as templates using standard procedures. Control in vitro transcripts have been synthesized on linearized plasmid DNAs of the precursor plasmids of pVE293 and pVE294 without the pds fragment insert and on linearized plasmid DNA of the infective clone STMV-10 of the wild type.

25

The in vitro transcripts have been mechanically inoculated onto leaves of three months old *Petunia hybrida* V26 plants together with the TMV-U2 helper virus.

30

The infected plants have been continuously scored for pds inactivation, which resulted in a phenotype showing leaf bleaching. The infection of petunia plants with TMV/STMV caused the occurrence of crinkled leaves starting from the infected branches but quickly progressing systemically throughout the plant. The

plants did survive the infections and often showed recovery phenotypes with almost no symptoms.

Only the plants, which have been infected with the hybrid vectors pVE293 and pVE294 carrying the pds fragment, showed in addition to the virus symptoms, additional phenotypic changes. In case of TMV/pVE293 infections, the infected branches developed young leaves with bleached sectors around the veins and at the tip of the leaves. This bleaching was progressive and independent from the presence of virus symptoms. Some of the leaves were fully bleached and progressively changed into almost white color. This phenotype was restricted to leaves of the infected branches. In case of TMV/pVE294 infections a similar bleaching of young leaves was observed for all developing branches but not for branches carrying already terminal flower buds. The bleaching started again around the veins and leaf tips but proceeded quickly producing variable leaf variegation patterns. This phenotype has never been observed in control inoculations with wild type viruses or deletion mutants. Therefore this phenotype indicates the functional knockout of pds in *Petunia hybrida* plants after infection with pVE293 and pVE294 viral hybrid vectors.

In RNA gel blot analyses performed with total RNA preparations of green and bleached spots of upper leaves of plants showing a FKO phenotype no chimeric STMV virus RNA and only low levels of TMV helper virus RNA could be detected. This indicates that virus induced gene silencing of pds in a specific tissue does not need to be accompanied by high levels of virus RNA.

REFERENCES

Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, Current Protocols, USA.

Baulcombe (1996) *Plant Cell* 8: 1833-1844

5 Baulcombe *et al.* (1998) JIC &SL Annual Report 1996/1997

Bringloe *et al.* (1998) 79: 1539-1546

Chapman (1991) PhD dissertation, University of Cambridge, UK

Coutts *et al.* (1991) *J. Gen. Virology* 72: 1521-1529

Danthinne *et al.* (1991) *Virology* 185: 605-614

10 Depicker and Van Montagu (1997) *Curr. Opin. Cell. Biol.* 9: 373-382

English *et al.* (1996) *Plant Cell* 8, 179-188

Hamilton *et al.* (1998) *The Plant Journal* 15(6): 737-746

Kempin *et al.* (1997) *Nature* 389: 802-803

Kumagai *et al.* (1995) *Proc. Natl. Acad. Sci USA* 92: 1679-1683

15 Meulewaeter *et al.* (1990) *Virology* 177:699-709

Pereira and Aerts (1998) *Methods in Molecular Biology* 82 Eds. Martinez-Zapatar and Salinas, Humana Press, NJ

R.D.D. Croy (1993) *Plant Molecular Biology Labfax* jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

20 Prahasar *et al.* (1996) *Proc. Natl. Acad. Sci USA* 93: 659-663

Ruiz *et al.* (1998) *The Plant Cell* 10: 937-946

Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, NY

Stam *et al.* (1997) *Ann. Botan.* 79:3-12

25 Takayuki *et al.* (1995) *The Plant Journal* 8(5):771-776

Walkey (1985) *Applied Virology*, William Heinemann Ltd, London

Waterhouse *et al.* (1998) *Proc. Natl. Acad. Sci USA* 95: 13959-13964

Wellink *et al.* (1998) Abstract presented at the Joint Meeting of Arbeitskreis Virologie and Nederlandse Kring voor Plantenvirologie in Wageningen, The Netherlands, November 12 and 13, 1998.

30

Wilbur and Lipmann (1983) *Proc. Nat. Acad. Sci. USA* 80: 726

Ysebaert *et al.* (1980) *J. Mol. Biol.* 143: 273-287

Claims

1. A method for isolating genes involved in the determination of a trait or phenotype of a plant species, said method comprising
 - a.) Identifying a set of nucleic acid sequences of genes, whose expression is correlated with a trait of interest.
 - b.) creating a library of gene silencing constructs in a viral RNA vector, said viral RNA vector being capable of replication inside plant cells and said gene silencing constructs being targeted to the nucleotide sequence of said nucleic acid sequences;
 - c.) infecting a collection of individual plants of said plant species with said library of gene silencing constructs, whereby each plant is infected with at least one member of said library;
 - d.) identifying a plant wherein said trait or phenotype is altered, using an assay adapted to said trait or phenotype;
 - e.) isolating said gene involved in the determination of said trait or phenotype in said plant species, from said library based on the nucleotide sequence to which said gene silencing construct in said identified plant was targeted.
2. The method of claim 1, wherein said viral RNA vector is capable of autonomous replication inside plant cells.
3. The method of claim 2, wherein said viral RNA vector is derived from cowpea mosaic virus.
4. The method of claim 1, wherein said viral RNA vector is capable of replication inside plant cells when the required factors are supplemented *in trans*.

5. The method of claim 4, wherein said viral RNA vector is derived from a satellite RNA virus and said factors are supplemented by infection with a helper virus or helper virus RNA.

5 6. The method of claim 5, wherein said satellite RNA virus is satellite tobacco mosaic virus.

7. The method of claim 6, wherein said viral RNA vector comprises an origin of assembly of tobacco mosaic virus.

10

8. The method of claim 7, wherein said helper virus is tobacco mosaic virus.

15

9. The method of claim 5, wherein said satellite RNA virus is satellite tobacco necrosis virus.

10. The method of claim 9, wherein said viral RNA vector comprises an origin of assembly of tobacco mosaic virus.

20

11. The method of claim 10, wherein said helper virus is derived from tobacco necrosis virus and comprises a coat protein gene of tobacco mosaic virus.

25

12. The method of claim 11, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said helper virus is derived from TNV-A.

13. The method of claim 11, wherein said satellite RNA virus is STNV-C and said helper virus is derived from TNV-D.

30

14. The method of claim 1, wherein said gene-silencing constructs comprise antisense RNA.

15. The method of claim 1, wherein said gene-silencing constructs comprise sense RNA.

16. The method of claim 1, wherein said gene-silencing constructs comprise an inverted repeat.

17. The method of claim 1, wherein said gene-silencing constructs comprise complementary stretch of at least 50 nucleotides of sense and antisense RNA.

18. The method of claim 17, wherein said gene-silencing constructs comprise complementary stretch of at least 100 nucleotides of sense and antisense RNA.

19. The method of claim 17, wherein said gene-silencing constructs comprise at least two copies of part of the nucleotide sequences of said collection of nucleic acids, said copies being in inverted repeat.

20. A method for the isolation of a nucleic acid with a specific function from a collection of nucleic acids, said collection of nucleic acids being characterized in that variation in the expression pattern of said nucleic acids is correlated with variation in a trait/phenotype of a plant harboring said nucleic acids, said method comprising the steps of

a.) creating a library of gene silencing constructs in a viral RNA vector, said viral RNA vector being capable of replication inside, and said gene silencing constructs being targeted/adapted to the nucleotide sequence of said nucleic acids;

b.) infecting a collection of plants with said library of gene silencing constructs, whereby each plant is infected with at least one member of said library;

c.) identifying plants with altered trait or phenotype using an assay adapted to said trait or phenotype.

21. The method of claim 20, further comprising the step of isolating said
5 nucleic acid with said specific function from said identified plant with altered trait or phenotype.

22. A method for determining the function encoded by a nucleic acid comprising a known nucleotide sequence in a plant, said method comprising

10 a.) providing a viral RNA vector, said viral RNA vector being derived from a satellite RNA virus, comprising a gene-silencing construct targeted to a gene comprising said known nucleotide sequence;

b.) infecting said plant with said viral RNA vector and a
15 corresponding helper virus;

c.) identifying an altered trait or phenotype of said co-infected plant.

23. A method for isolating essential genes in a plant, comprising

a.) creating a library of random gene-silencing constructs for said
20 plant comprised within a viral RNA vector, said viral RNA vector being derived from a satellite RNA virus;

b.) infecting a plant with at least one member of said library and with a corresponding helper virus; and

c.) identifying plants developing a gene-silencing-construct-
25 associated phenotype, preferably chlorosis or necrosis.

24. The method according to claim 23, further comprising the step of isolating the viral RNA vector from the tissue exhibiting the phenotype.

30 25. The method of claim 23, wherein said library is created by cloning random DNA fragments of said plant in a cDNA copy of the viral RNA vector.

26. The method of claim 23, wherein said library is created by cloning random cDNA fragments of said plant in a cDNA copy of the viral RNA vector.

27. The method of claim 23, wherein said library is created by cloning
5 random duplicated cDNA fragments in inverted repeat.

28. The method of claim 23, wherein said viral RNA vector is derived from STMV and comprises an origin of assembly of tobacco mosaic virus, and wherein said helper virus is tobacco mosaic virus.

10

29. The method of claim 23, wherein said viral RNA vector is derived from satellite tobacco necrosis virus and comprises an origin of assembly of tobacco mosaic virus and wherein said helper virus is derived from tobacco necrosis virus and comprises a coat protein gene of tobacco mosaic virus.

15

30. The method of claim 29, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said helper virus is derived from TNV-A.

20

31. The method of claim 29, wherein said satellite RNA virus is STNV-C and said helper virus is derived from TNV-D.

32. A method for the introduction of inhibitory RNA in the cytoplasm of plant cells, said method comprising:

25

a.) introducing into said plant cell, a viral RNA vector comprising said inhibitory RNA or comprising a chimeric nucleic acid which when transcribed yields said inhibitory RNA, wherein said viral RNA vector is derived from a satellite RNA virus; and

b.) introducing a corresponding helper virus into said plant cell.

30

33. The method of claim 32, wherein said inhibitory RNA comprises sense RNA.

34. The method of claim 32, wherein said inhibitory RNA comprises antisense RNA.

5 35. The method of claim 32, wherein said inhibitory RNA comprises an inverted repeat.

36. The method of claim 32, wherein said inhibitory RNA comprises complementary stretch of at least 50 nucleotides of sense and antisense
10 RNA.

37. The method of claim 36, wherein said inhibitory RNA comprises complementary stretch of at least 100 nucleotides of sense and antisense
15 RNA.

38. The method of claim 32, wherein said viral RNA vector is derived from STMV and comprises an origin of assembly of tobacco mosaic virus, and wherein said helper virus is tobacco mosaic virus.

20 39. The method of claim 32, wherein said viral RNA vector is derived from satellite tobacco necrosis virus and comprises an origin of assembly of tobacco mosaic virus and wherein said helper virus is derived from tobacco necrosis virus and comprises a coat protein gene of tobacco mosaic.

25 40. The method of claim 39, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said helper virus is derived from TNV-A.

41. The method of claim 39, wherein said satellite RNA virus is STNV-C
30 and said helper virus is derived from TNV-D.

42. The method of any one of claims 32 to 41, wherein said plant is selected from *Nicotiana* spp, *Oryza sativa*, *Zea Mays*, *Brassica* spp. , *Gossypum* spp., *Triticum* spp., *Arabidopsis* spp. or *Petunia* spp.

5

43. A kit for introduction of inhibitory RNA in the cytoplasm of a plant cell, said kit comprising

- a.) a viral RNA vector derived from a satellite RNA virus, said vector comprising a chimeric nucleic acid which when transcribed yields said inhibitory RNA or which comprises said inhibitory RNA; and
- b.) a corresponding helper virus.

10

44. The kit of claim 43, wherein said inhibitory RNA comprises sense RNA.

15

45. The kit of claim 43, wherein said inhibitory RNA comprises antisense RNA.

46. The kit of claim 43, wherein said inhibitory RNA comprises an inverted repeat.

20

47. The kit of claim 43, wherein said inhibitory RNA comprises complementary stretch of at least 50 nucleotides of sense and antisense RNA.

25

48. The kit of claim 43, wherein said inhibitory RNA comprises complementary stretch of at least 100 nucleotides of sense and antisense RNA.

30

49. The kit of claim 43, wherein said viral RNA vector is derived from STMV and comprises an origin of assembly of tobacco mosaic virus, and wherein said corresponding helper virus is tobacco mosaic virus.

50. The kit of claim 43, wherein said viral RNA vector is derived from satellite tobacco necrosis virus and comprises an origin of assembly of tobacco mosaic virus and wherein said corresponding helper virus is derived from tobacco necrosis virus and comprises the coat protein gene of tobacco mosaic virus .


51. The kit of claim 50, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said corresponding helper virus is derived from TNV-A.

52. The kit of claim 50, wherein said satellite RNA virus is STNV-C and said corresponding helper virus is derived from TNV-D.


Abstract

- 5 The invention provides methods and means for the identification of genes involved in the determination of a plant trait or for the identification encoded by a nucleic acid comprising a determined nucleotide sequence. The invention also provides kits comprising viral RNA vectors derived from satellite viruses and corresponding helper viruses for the introduction of
- 10 inhibitory RNA into plant cells and plants.

1. Tobacco mosaic virus




2. Tobacco necrosis virus



1/1

3. Satellite tobacco mosaic virus



4. Satellite tobacco necrosis virus



COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY
(Includes Reference to Provisional and PCT International Applications)

Attorney's Docket No.

021565-075

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHODS AND MEANS FOR DELIVERING INHIBITORY RNA TO PLANTS AND APPLICATIONS THEREOF

the specification of which (check only one item below):

☒ is attached hereto.

☐ was filed as United States application

Number _____
on _____
and was amended
on _____ (if applicable).

☐ was filed as PCT international application

Number _____
on _____
and was amended
on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(e) of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. §119:

COUNTRY (if PCT, indicate "PCT")	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 U.S.C. §119
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below.

(Application Number)

(Filing Date)

(Application Number)

(Filing Date)

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (CONT'D)
(Includes Reference to Provisional and PCT International Applications)

Attorney's Docket No.

021565-075

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose to the Office all information known to me to be material to the patentability as defined in Title 37, Code of Federal Regulations §1.56, which became available between the filing date of the prior application(s) and the national or PCT international filing date of this application:

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. §120:

U.S. APPLICATIONS		STATUS (check one)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	PATENTED	PENDING	ABANDONED
09/294,022	April 20, 1999		X	
PCT APPLICATIONS DESIGNATING THE U.S.				
PCT APPLICATION NO.	PCT FILING DATE	U.S. APPLICATION NUMBERS ASSIGNED (if any)		

I hereby appoint the following attorneys and agent(s) to prosecute said application and to transact all business in the Patent and Trademark Office connected therewith and to file, prosecute and to transact all business in connection with international applications directed to said invention:

William L. Mathis	17,337	R. Danny Huntington	27,903	Gerald F. Swiss	30,113
Robert S. Swecker	19,885	Eric H. Weisblatt	30,505	Michael J. Ure	33,089
Platon N. Mandros	22,124	James W. Peterson	26,057	Charles F. Wieland III	33,096
Benton S. Duffett, Jr.	22,030	Teresa Stanek Rea	30,427	Bruce T. Wieder	33,815
Norman H. Stepno	22,716	Robert E. Krebs	25,885	Todd R. Walters	34,040
Ronald L. Grudziecki	24,970	William C. Rowland	30,888	Ronni S. Jillions	31,979
Frederick G. Michaud, Jr.	26,003	T. Gene Dillahunt	25,423	Harold R. Brown III	36,341
Alan E. Kopecki	25,813	Patrick C. Keane	32,858	Allen R. Baum	36,086
Regis E. Slutter	26,999	Bruce J. Boggs, Jr.	32,344	Steven M. du Bois	35,023
Samuel C. Miller, III	27,360	William H. Benz	25,952	Brian P. O'Shaughnessy	32,747
Robert G. Mukai	28,531	Peter K. Skiff	31,917	Kenneth B. Leffler	36,075
George A. Hovanec, Jr.	28,223	Richard J. McGrath	29,195	Fred W. Hathaway	32,236
James A. LaBarre	28,632	Matthew L. Schneider	32,814		
E. Joseph Gess	28,510	Michael G. Savage	32,596		



21839

and: Malcolm K. McGowan, Ph.D., Reg. No. 39,300

Address all correspondence to:



21839

R. Danny Huntington, Esq.
BURNS, DOANE, SWECKER & MATHIS, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404

Address all telephone calls to: Malcolm K. McGowan, Ph.D. at (703) 836-6620.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (CONT'D) (Includes Reference to Provisional and PCT International Applications)	Attorney's Docket No. 021565-075
---	-------------------------------------

FULL NAME OF SOLE OR FIRST INVENTOR	SIGNATURE	DATE
Frank Meulewaeter		
RESIDENCE	CITIZENSHIP	
Eksaarde, Belgium	Belgian	
POST OFFICE ADDRESS		
Weehaagstraat 73, 9160 Eksaarde, Belgium		
FULL NAME OF SECOND JOINT INVENTOR, IF ANY	SIGNATURE	DATE
Marc Cornelissen		
RESIDENCE	CITIZENSHIP	
Heusden, Belgium	Dutch	
POST OFFICE ADDRESS		
Ellebogten 38, 9070 Heusden, Belgium		
FULL NAME OF THIRD JOINT INVENTOR, IF ANY	SIGNATURE	DATE
John Jacobs		
RESIDENCE	CITIZENSHIP	
Roosendaal, Netherlands	Dutch	
POST OFFICE ADDRESS		
Dilleberg 20, 4708 DL Roosendaal, The Netherlands		
FULL NAME OF FOURTH JOINT INVENTOR, IF ANY	SIGNATURE	DATE
Gerben Van Eldik		
RESIDENCE	CITIZENSHIP	
Gent, Belgium	Dutch	
POST OFFICE ADDRESS		
Nekkersberglaan 41, 9000 Gent, Belgium		
FULL NAME OF FIFTH JOINT INVENTOR, IF ANY	SIGNATURE	DATE
Michael Metzlauff		
RESIDENCE	CITIZENSHIP	
Tervuren, Belgium	German	
POST OFFICE ADDRESS		
Irislaan 26, 3080 Tervuren, Belgium		
FULL NAME OF SIXTH JOINT INVENTOR, IF ANY	SIGNATURE	DATE
RESIDENCE	CITIZENSHIP	
POST OFFICE ADDRESS		
FULL NAME OF SEVENTH JOINT INVENTOR, IF ANY	SIGNATURE	DATE
RESIDENCE	CITIZENSHIP	
POST OFFICE ADDRESS		
FULL NAME OF EIGHTH JOINT INVENTOR, IF ANY	SIGNATURE	DATE
RESIDENCE	CITIZENSHIP	
POST OFFICE ADDRESS		

SEQUENCE LISTING

<110> Meulewater, Frank
Cornelissen, Marc
Van Eldik, Gerben
Jacobs, John

<120> Methods and means for delivering inhibitory RNA to
plants and applications thereof

<130> FKOSAT

<140>

<141>

<160> 15

<170> PatentIn Ver. 2.0

<210> 1

<211> 3684

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:cDNA copy of
the nucleotide sequence of the genome of TNV-A

<400> 1

```

agtattcata ccaagaatac caaatagggtg caaggcctta ctcagctaaa gagtctaaaa 60
tgagagctacc aaaccaacac aagcaaacgg ccgccgagggg tttcgatatct ttccctaaact 120
ggctatgcaa cccatggaga cgacagcgaa cagtcaacgc tgcagttgag ttccaaaaag 180
atcttctcgc cattgaggat tccgagcatt tggatgacat caatgagtgt ttcgaggagt 240
ctgctggggc acaatctcag cgaactaagg ttgtcgccga cggagcatat gccccgcaa 300
aatccaacag gacccgccga gttcgtaagc agaagaagca caagtttgta aaatatcttg 360
tcaacgaagc tcgtgccgag tttggattgc ccaaaccaac tgaggcaaac agacttatgg 420
tccaacattt cttgctcaga gtgtgcaagg attggggcgt tgttactgcc cacgtacacg 480
gcaatgttgc actagctttg cactgggtgt tcatcccaac ggaagatgat ctgctatcac 540
gagcattgat gaacacacat gctactagag ccgctgtacg aggcattggac aatgtccaag 600
gggaggggtg gtggaacaat aggttgggga ttggggggcca ggtcggactg gccttcgggt 660
ccaaataggg gtgccttgaa aggaggccag gattctccac gtccgtttcg cgtggggaac 720
atcctgatct ggtggtcata ccatcagggc gccctgagaa acagcgtcag ttgttacgct 780
atagtggat aggcggccat ttattaatcg gcatccacaa caactctctt tccaacctgc 840
gtaggggctt gatggaaaga gtattctatg tcgagggggc caatgggctt caagacgccc 900
ctaagcccg caagggagct ttcgaaccc ttgataagtt tcgtgatctc tatactaaaa 960
atagtgtggc tcatacccct gtaactagt aacaattcct aatgaattac acgggcagga 1020
aactgactat ttacagagag gcggttgata gtttgtcgca tcaaccctt agctcacgag 1080
atgcgaaact aaagacattc gtgaaggccg aaaaattaaa tctttctaag aagcctgacc 1140

```

ctgctcccag	ggatcatccaa	cctagatcgc	ctcgggtataa	cgtttgtttg	ggcagggtacc	1200
tccgacatta	tgagcatcac	gcgttttaaaa	ccattgccaa	gtgctttggg	gaaatcacgg	1260
tcttcaaagg	gtttactctg	gagcaacaag	gggaaatcat	gcgctcgaag	tggaataaat	1320
atgttaatcc	cgtcgcagtc	ggactcgacg	ccagtcgttt	cgaccaacac	gtgtctgttg	1380
aagcactcga	gtatgagcat	gaattttacc	tcagagacta	cccaaataat	aaacagctaa	1440
aatggctgct	aaagcagcaa	ttgtgcaacg	taggaacggc	attcgccagt	gacggcatta	1500
taaaatacaa	gaagaagggt	tgtagaatga	gcggagacat	gaacacgagt	ttgggcaact	1560
gcatttctaat	gtgcgccatg	gtctacgggt	tgaaagaaca	cttaaacatc	aatttggtccc	1620
ttgcaaataa	tggggatgac	tgcgtcattg	tctgtgagaa	agcggattta	aagaaattga	1680
caagcagcat	cgagccatat	ttcaagcagt	ttggattcaa	gatggaagtg	gaaaaacccg	1740
tggatatatt	tgagcgcata	gaattttgcc	aaaccaaac	tgtgttcgat	ggatcccagt	1800
acatcatggt	acgcaaacct	tctgtggtta	catctaaaga	cgtcactagc	cttatcccat	1860
gtcaaacgaa	agcacaatac	gcagaatggc	tgcaagctgt	aggtgagtgt	ggcatgagca	1920
ttaacgggtg	gattcctgtc	atgcagaatt	tctacaaaa	gctccaaact	ggcatccgcc	1980
gcacaaaatt	caccaagacc	ggcgagttcc	agacgaacgg	attgggggat	cactctagat	2040
atatgcatag	agtggcccg	gttccttcgc	ctgaaacccg	tttatccttc	tatctagctt	2100
tcggtatcac	accagacctc	caagaagcat	tggagatctt	ctatgatacc	cacaggcttg	2160
agttggatga	tggtatccca	actgatacct	accaagtgtc	aggagagcat	ttgatcaatg	2220
gattaccaa	ctgatgtaac	ggaggacaat	gtgcaaatac	gcggtcgggc	taggagcggt	2280
gagggtaaga	aacacaatgg	ttcgggatta	actggcggtta	agcgtcacgc	ggtgagcgaa	2340
acatctcaga	aatcacagca	aggtactggc	aatggaacta	tgaccaatat	agccgaagaa	2400
cagaccatta	ccgtgacata	caactttaac	ttttaagtta	tgggtgcgtg	tcgctgttgt	2460
gatacttcac	cagggtattac	actattccct	tactttgcaa	ttctcatcct	tatattggca	2520
atacttggtg	tagggactcc	caatcaacaa	tatcaccatt	ctccaagcac	ttacgagtac	2580
aagactcaac	acatttcgat	cgcaaaatag	acatggcagg	aaagaagaac	aacaacaacg	2640
gtcagtatat	aatactgcgt	actccagagc	aacaggtgga	gatagaccag	cgcaacgccc	2700
gtcgtgctca	aatgggtcgc	atgaagaagg	ctagacagcc	cgttcagcga	tacttacagc	2760
aacacgggtt	gcgaaacgga	ttgtccggta	gagggggcta	catagtggct	cccacctccg	2820
gggggggtgt	cactcgaccc	atagtgccga	aattctccaa	caggggagat	tccactatag	2880
tccgtaacac	tgagattttg	aacaacccaa	tcttagcggc	gctaggcgca	ttcaatacaa	2940
caaactccgc	actgattgca	gcagcaccat	catggctggc	tagcatcgct	gatctttaca	3000
gtaaatatag	atggctctca	tgtgagatca	tctacattcc	aaaatgcccc	accaccacca	3060
gtggatcaat	tgccatggct	ttcacatacg	acagaaatga	cgctgcaccc	accgcaaggg	3120
ctcagctgtc	acaatcttac	aaggccatca	atcttccacc	gtatgcggga	tacgacggag	3180
cagcatattt	gaattcgaac	caggagagctg	ggtcagccat	cgccgttcaa	cttgatgtta	3240
ccaagttgga	caagccatgg	taccccaacta	tctcctctgc	cggcttcggg	gcgctcagcg	3300
tcctcgatca	gaaccaattc	tgccccgcgt	cccttggtgt	cgctagcgat	gggggacccg	3360
ctactgctac	tccagcaggg	gaccttttca	tcaagtacgt	gattgagttc	attgaaccaa	3420
tcaaccacac	aatgaacgtc	tagttctttg	tactgttaact	tggctaattg	ctaaggtgga	3480
gtcacaccat	tggagacgga	gacggatcct	gggaaacagg	cttgacgggc	gggggggtgt	3540
gccccgacg	acgcatcact	ccggatacca	atgggtacacc	actatggcag	ggtctgcca	3600
ggtcttgtgc	accaagaacc	cctggaaacg	ggggggaggg	gggtagcaca	tatcatccag	3660
attgaggggc	ctttgcccc	cccc				3684

<210> 2

<211> 6395

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of TMV-U1

<400> 2

gtatTTTTac aacaattacc aacaacaaca aacaacaaac aacattacaa ttactatttta 60
caattacaat ggcatacaca cagacagcta ccacatcagc tttgctggac actgtccgag 120
gaaacaactc cttgggtcaat gatctagcaa agcgtcgtct ttacgacaca gcggttgaag 180
agtttaacgc tcgtgaccgc aggcccaagg tgaacttttc aaaagtaata agcgaggagc 240
agacgcttat tgctacccgg gcgtatccag aattccaaat tacattttat aacacgcaaa 300
atgccgtgca ttcgcttgca ggtggattgc gatctttaga actggaatat ctgatgatgc 360
aaattcccta cggatcattg acttatgaca taggcgggaa ttttgcacgc catctgttca 420
agggacgagc atatgtacac tgctgcatgc ccaacctgga cgttcgagac atcatgcggc 480
acgaaggcca gaaagacagt attgaactat acctttctag gctagagaga ggggggaaaa 540
cagtcccaa cttccaaaag gaagcatttg acagatacgc agaaattcct gaagacgctg 600
tctgtcacia tactttccag acaatgcgac atcagccgat gcagcaatca ggcagagtgt 660
atgccattgc gctacacagc atatatgaca taccagccga tgagtccggg gcggcactct 720
tgaggaaaaa tgtccatacg tgctatgccg ctttccactt ctctgagaac ctgcttcttg 780
aagattcata cgtcaatttg gacgaaatca acgcgtgttt ttcgcgcgat ggagacaagt 840
tgaccttttc ttttgcacga gagagtactc ttaattattg tcatagttat tctaataattc 900
ttaagtatgt gtgcaaaact tacttcccgg cctctaatag agaggtttac atgaaggagt 960
ttttagtcac cagagttaat acctggtttt gtaagttttc tagaatagat acttttcttt 1020
tgtacaaaag tgtggcccat aaaagtgtag atagtgcgca gttttatact gcaatggaag 1080
acgcatggca ttacaaaaag actcttgcaa tgtgcaacag cgagagaatc ctcttgagg 1140
attcatcatc agtcaattac tgggttccca aaatgaggga tatggtcac gtaccattat 1200
tcgacatttc tttgggagact agtaagagga cgcgcaagga agtcttagtg tccaaggatt 1260
tcgtgtttac agtgcttaac cacattcgaa cataccaggc gaaagctctt acatacgcaa 1320
atgttttgtc ctttgcgaa tcgattcgat cgagggtaat cattaacggt gtgacagcga 1380
ggtccgaatg ggatgtggac aaatctttgt tacaatcctt gtccatgacg ttttacctgc 1440
atactaagct tgccgttcta aaggatgact tactgattag caagtttagt ctccggttcga 1500
aaacggtgtg ccagcatgtg tgggatgaga tttcgctggc gtttgggaac gcatttccct 1560
ccgtgaaaga gaggtctctg aacaggaaac ttatcagagt ggcaggcgac gcattagaga 1620
tcagggtgcc tgatctatat gtgaccttcc acgacagatt agtgactgag tacaaggcct 1680
ctgtggacat gcctgcgctt gacattagga agaagatgga agaaacggaa gtgatgtaca 1740
atgcactttc agagttatcg gtgttaaggg agtctgacaa attcgatgtt gatgtttttt 1800
cccagatgtg ccaatctttg gaagttgacc caatgacggc agcgaagggt atagtcgcgg 1860
tcatgagcaa tgagagcggc ctgactctca catttgaaac acctactgag gcgaatgttg 1920
cgctagcttt acaggatcaa gagaaggctt cagaagggtgc tttggtagt acctcaagag 1980
aagttgaaga accgtccatg aagggttcga tggccagagg agagttacaa ttagctgggtc 2040
ttgctggaga tcatccggag tcgtcctatt ctaagaacga ggagatagag tcttttagagc 2100
agtttcatat ggcaacggca gattcggtta ttcgtaagca gatgagctcg attgtgtaca 2160
cgggtccgat taaagttcag caaatgaaaa actttatcga tagcctggta gcactactat 2220
ctgctgcggg gtcgaatctc gtcaagatcc tcaagatac agctgctatt gaccttgaaa 2280
cccgtcaaaa gtttggagtc ttggatgttg catctaggaa gtggttaatc aaaccaacgg 2340
ccaagagtca tgcatggggg gttgttgaaa ccacgcgag gaagtatcat gtggcgcttt 2400
tggaatatga tgagcagggt gtggtgacat gcgatgattg gagaagagta gctgtcagct 2460
ctgagtctgt tgtttattcc gacatggcga aactcagaac tctgcgcaga ctgcttcgaa 2520

acggagaacc gcatgtcagt agcgcaaagg ttgttcttgt ggacggagtt ccgggctgtg 2580
 ggaaaaccaa agaaattcct tccaggggta attttgatga agatctaatt ttagtacctg 2640
 ggaagcaagc cgcggaatg atcagaagac gtgcgaattc ctcagggatt attgtggcca 2700
 cgaaggacaa cggttaaaacc gttgattcct tcatgatgaa ttttgggaaa agcacacgct 2760
 gtcagttcaa gaggttattc attgatgaag ggttgatggt gcatactggt tgtgttaatt 2820
 ttcttgtggc gatgtcattg tgcgaaattg catatgttta cggagacaca cagcagattc 2880
 catacatcaa tagagtttca ggattcccgt accccgccca ttttgccaaa ttggaagttg 2940
 acgaggtgga gacacgcaga actactctcc gttgtccagc cgatgtcaca cattatctga 3000
 acaggagata tgagggcttt gtcatgagca cttcttcggt taaaaagtct gtttcgcagg 3060
 agatggtcgg cgagcgccgc gtgatcaatc cgatctcaaa acccttgcat ggcaagatcc 3120
 tgacttttac ccaatcggat aaagaagctc tgctttcaag agggatttca gatgttcaca 3180
 ctgtgcatga agtgcaaggc gagacatact ctgatgtttc actagttagg ttaaccctta 3240
 caccagtctc catcattgca ggagacagcc cacatgtttt ggtcgcattg tcaaggcaca 3300
 cctgttcgct caagtactac actgttggtt tggatccttt agttagtatc attagagatc 3360
 tagagaaact tagctcgtac ttgttagata tgtataaggt cgatgcagga acacaatagc 3420
 aattacagat tgactcgggt ttcaaagggt ccaatctttt tgttgcagcg ccaaagactg 3480
 gtgatatttc tgatatgcag ttttactatg ataagtgtct cccaggcaac agcaccatga 3540
 tgaataatth tgatgctggt accatgaggt tgactgacat ttcattgaat gtcaaagatt 3600
 gcatattgga tatgtctaag tctgttgctg cgcctaagga tcaaatacaa ccactaatac 3660
 ctatggtacg aacggcggca gaaatgccac gccagactgg actattggaa aatttagtgg 3720
 cgatgattaa aaggaacttt aacgcacccg agttgtctgg catcattgat attgaaaata 3780
 ctgcatcttt agttgtagat aagttttttg atagttatth gcttaaagaa aaaagaaaac 3840
 caaataaaaa tgtttctttg ttcagtagag agtctctcaa tagatggtta gaaaagcagg 3900
 aacaggtaac aataggccag ctgcgagatt ttgattttgt agatttgcca gcagttgatc 3960
 agtacagaca catgattaaa gcacaacca agcaaaaatt ggacacttca atccaaacgg 4020
 agtaccgggc tttgcagacg attgtgtacc attcaaaaaa gatcaatgca atatttggcc 4080
 cgttgtttag tgagcttact aggcaattac tggacagtgt tgattcgagc agatttttgt 4140
 ttttcacaag aaagacacca gcgcagattg aggatttctt cggagatctc gacagtcag 4200
 tgccgatgga tgtcttggag ctggatatat caaaatacga caaatctcag aatgaattcc 4260
 actgtgcagt agaatacgag atctggcgaa gattgggttt tgaagacttc ttgggagaag 4320
 tttgaaaaca agggcataga aagaccaccc tcaaggatta taccgcagggt ataaaaactt 4380
 gcatctggta tcaaagaaag agcggggacg tcacgacggt cattggaaac actgtgatca 4440
 ttgtcgcag tttggcctcg atgcttccga tggagaaaat aatcaaagga gccttttgcg 4500
 gtgacgatag tctgctgtac tttccaaagg gttgtgagtt tccggatgtg caacactccg 4560
 cgaatcttat gtggaattht gaagcaaaac tgtttaaaaa acagtatgga tacttttgcg 4620
 gaagatatgt aatacatcac gacagaggat gcattgtgta ttacgatccc cttaaagtga 4680
 tctcgaaact tgggtgctaaa cacatcaagg attgggaaca cttggaggag ttcagaagg 4740
 ctctttgtga tgttgctggt tegtgtgaaca attgtgcgta ttacacacag ttggacgacg 4800
 ctgtatggga ggttcataag accgcccctc caggttcgtt tgtttataaa agtctggtga 4860
 agtatttgtc tgataaagtt ctttttagaa gtttgthtat agatggctct agttgttaaa 4920
 ggaaaagtga atatcaatga gtttatcgac ctgacaaaaa tggagaagat cttaccgtcg 4980
 atgtttaccc ctgtaaagag tgttatgtgt tccaaagttg ataaaaataat ggttcatgag 5040
 aatgagtcac tgtcagagggt gaaccttctt aaaggagtta agcttattga tagtggtac 5100
 gtctgttttag ccggttttgt cgtcacgggc gagtggaaact tgcctgacaa ttgcagagga 5160
 ggtgtgagcg tgtgtctggt ggacaaaagg atggaaagag ccgacgaggc cactctcgga 5220
 tcttactaca cagcagctgc aaagaaaaga tttcagttca aggtcgttcc caattatgct 5280
 ataaccaccc aggacgcgat gaaaaacgtc tggcaagttt tagttaatat tagaaatgtg 5340
 aagatgtcag cgggtttctg tccgctttct ctggagtttg tgtcgggtgtg tattgtttat 5400


```

agaaataata taaaattagg tttgagagag aagattacaa acgtgagaga cggagggccc 5460
atggaactta cagaagaagt cgttgatgag ttcattggaag atgtccctat gtcatcagg 5520
cttgcaaagt ttcgatctcg aaccggaaaa aagagtgatg tccgcaaagg gaaaaatagt 5580
agtaatgatc ggtcagtgcc gaacaagaac tatagaaatg ttaaggattt tggaggaatg 5640
agtttttaaaa agaataattt aatcgatgat gattcggagg ctactgtcgc cgaatcggat 5700
tcgttttaaaa tatgtcttac agtatcacta ctccatctca gttcgtgttc ttgtcatcag 5760
cgtggggccga cccaatagag ttaattaatt tatgtactaa tgccttagga aatcagtttc 5820
aaacacaaca agctcgaact gtcgttcāaa gacaattcag tgaggtgtgg aaaccttcac 5880
cacaagtaac tgtaggttc cctgacagtg actttaaggt gtacaggtac aatgcggtat 5940
tagaccgct agtcacagca ctgttaggtg cattcgacac tagaaataga ataatagaag 6000
ttgaaaatca ggcgaacccc acgactgccg aaacgttaga tgctactcgt agagtagacg 6060
acgcaacggt ggccataagg agcgcgataa ataatttaat agtagaattg atcagaggaa 6120
ccggatctta taatcggagc tctttcgaga gctcttctgg tttggtttg acctctggtc 6180
ctgcaacttg aggtagtcaa gatgcataat aaataacgga ttgtgtccgt aatcacacgt 6240
ggtgcgtacg ataacgcata gtgtttttcc ctccacttaa atcgaagggt tgtgtcttg 6300
atcgcgcggg tcaaattgat atggttcata tacatccgca ggcacgtaat aaagcgaggg 6360
gttcgaatcc ccccgttacc cccggtaggg gccca 6395

```

<210> 3

<211> 1245

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of STNV-2

<400> 3

```

agtaaagaca ggaaacttta ccgactatca gaatgacaaa acgtcaaagc aaacaatcaa 60
accgcaagag cgttgcatca cagggtgcgtg gtattgttga gtcaatggct gagcagaagc 120
gatttgcttt tcttacgaac accaacacag tcaactacagc aggtaccgtg atcaacctga 180
gcaacaacat cgtgcaagga gatgaccttg ttaatcgcac cggagaccag attaagacca 240
tacaccagac tttattgact cgggtgtacag gaattaccaa cagccaaagc tttcggttca 300
tctggtttcg tgacaacacc aataggggga ctacaccggc tgtgactgag gtgttagaca 360
gtgctagtat aacatcccag tataacccca ctacgttcca gcaaaagagg ttcactgttt 420
tccaagattt catgttggtg acctctatag ttggacgtgt gattgtccat cggactgccg 480
ttgataagaa acggcgtgcg atattttaca acgggtgctgc ttctgtagcc gcgtcaaagt 540
gccccggtgc cacatttgta cttgtcattg gatcacatgc cactggacag tatgatgtga 600
cagccgagat tgtttatctg gacatgtaga ccatgggtcat gatgatgata gtgaaggacg 660
ctgaaagatg cgtagctacc ctctggtgc acttcctggt gcaaagcaga accaaaggg 720
acgggtgtac ggcgacagc agtcctgaac tagtaaatca ggaccgggag aaaaccagct 780
gacggctaaa tccattccca ctagtgtatt agtggaacga ggccccgcgt gaattgggg 840
ggctgcatgg ggtggaaaac catgtggtcg cagtcatctt tccatgcat tattgtctca 900
atacttggtg gcaacaatgc tgtaaatcaa cgtagcactc aacatcactt caaaaccccc 960
tccatgtcac aagaatcaag atgcatgtct gtgttttagcg gtatatattt tgcatccact 1020
tgatcgtgat tttgccctgg gcacctcgcg cggttggtac ccgcggagac tccccacagc 1080
aacatggcat taggcaggga taaggatatg tgactagaca aatgcgcgtg aagctggaaa 1140
gtccggttag cagtgggggt gtgcggaatg cagcctcaac aaggtatagc tgctgcatag 1200

```

gagatgtgaa cctttcaaac ttgaattcaa gtctcatgac tgccc

1245

<210> 4

<211> 1058

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of STMV

<400> 4

agtaaactta ccaatcaaaa gacctaacca acaggactgt cgtggtcatt tatgctgttg 60
ggggacatag ggggaaaaca tattgccttc ttctacaaga ggccttcagt cgccataatt 120
acttggcgcc caattttggg tttcagttgc tgtttccagc tatggggaga ggtaaggtta 180
aaccaaaccg taaatcgacg ggtgacaatt cgaatgttgt tactatgatt agagctggaa 240
gctatcctaa ggtcaatccg actccaacgt gggtcagagc catacctttc gaagtgtcag 300
ttcaatctgg tattgctttt aaagtaccgg tcgggtcact attttcggca aatttccgga 360
cagattcctt tacaagcgtc acagtgatga gtgtccgtgc ttggaccagc ttaacaccgc 420
cagtaaata gaacagtttt gtgaggctga agccattgtt caagactggg gactctactg 480
aggagttcga agggcgtgca tcaaacatca acacacgagc ttctgtaggg tacaggattc 540
caactaattt gcgtcagaat actgtggcag ccgacaatgt atgcgaagta agaagcaact 600
gtcgacaagt cgccttggtt atttcgtgtt gttttaactg aacctcgaca taagcctttt 660
ggatcgaagg ttaaacgatc cgctcctcgc ttgagcttga ggcggcgtat ctcttatgtc 720
aacagagaca ctttggtcta tggttgtata acaatagata gactcccgtt tgcaagatta 780
gggttaacag atcttgccgt tagtctgggt agcgcgtaac cggccttgat ttatggaata 840
gatccattgt ccaatggctt tgccaatgga acgccgacgt ggctgtataa tacgtcggtg 900
acaagtacga aatcttggtt gtgtttttcc ctccacttaa atcgaagggt tttgttttgg 960
tcttcccga cgcatacgtt agtgtgacta ccgttggtcg aaacaagtaa aacaggaagg 1020
gggttcgaat ccctccctaa ccgcgggtaa gcggccca 1058

<210> 5

<211> 6355

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of TMV-U2

<400> 5

gatgttttaa tagttttcga caacaacaat taaaacaaaa acaacatatt acaacaaca 60
aacaacaaca atggcacaca tacaatctat aattagcaac gcccttcttg aaagcgtgag 120
tggtaaaaac actctcggtt atgaccttgc aagaaggcgc atgtacgata cggccgtgga 180
agaatttaac gccgcgcacc gtagaccaa ggtcaacttt tccaaaacta ttagcgaaga 240
gcaaacgctt ctagtctcca acgcgtaccc ggagttccag attacctttt ataatactca 300
aaatgccgta cacagtttgg ctggaggttt gagagcatta gaattggaat atctgatgct 360
acaagttccc tatggatcgc cgacatatga tatagggtggg aactttgcag cacatttgtt 420

caaaggcagg gattacgtgc attgctgtat gcccaatctg gacatacgag atataatgag 480
 gcacgaagga caaaaggact caattgagat gtatttgtcc agattgtctc gttctaacaa 540
 ggtaattcct gagtttcaaa gggaggcttt taacaggat gcagaagctc ccaacgaagt 600
 ctgctgctct aaaacttttc aggattgtcg aatacatccg ccagagaata gtggtagaag 660
 atacgctgtt gctctgcaca gtttgtatga tttcctgtg catgagtttg gagctgcgtt 720
 aatatctaag aatatacatg tatgttatgc agcttccatt ttggcagaag cattattact 780
 agaccagacg gaggttacgc ttaatgaaat aggcgcaact ttcaaaagag aagggtgatga 840
 tgtttctttt ttctttgctg atgaaagtac tttaaattat agtcataaat acaaaaatat 900
 cttgcattat gtagttaaatt cttactttcc tgcttctagt agaatagttt actttaagga 960
 attttttagtc actaggggta atacttggtt ttgtaaattt accaaagtag atacctatat 1020
 tctgtacaag agtggttagac aagtaggggtg tgatagtgat cagttctatg aggcgatgga 1080
 agacgccttt gcttacaaga aaaccttggc catgttcaac actgaaagag caatcttttag 1140
 agacacggct tcgggttaact tttggttccc taagatgaag gacatggtga tagtaccgct 1200
 gtttgagggt tctattacca gcaaaaagat gacaaggagt gaggtcattg ttaatcgtga 1260
 cttcgtttac acagtgccta atcatatcag aacatatcaa gccaaagcgt taacttacca 1320
 gaacgtatta tctttcgtgg agtctataag atcccgcgtg ataatacaatg gtgttactgc 1380
 taggtctgaa tgggatgtag ataaagcaat tcttcaacct ttgtcaatga ctttcttctt 1440
 gcagactaag ctggctgcgc ttcaagacga tatagtaatg ggaaagtttc ggtgcttgga 1500
 taagaccact tctgaactta tttgggatga ggtgggcaaa ttttttgga acgttttccc 1560
 cactatcaaa gagagattgg tgagcaggaa aattctggat gtaagtgaga atgctctgaa 1620
 gatcaagatc ccagatctgt atgtcacatg gaaagacagg ttcgtagctg aatacaccaa 1680
 gtctgaggag ttaccgcac tagatatcaa gaaggactta gaagaagctg agcaaatgta 1740
 cgacgcgtta tcagaattat ctatccttaa ggtgctgat aatttcgata tcgcgaagtt 1800
 caaagacatg tgcaaggctt tagatgttag tctgatgtg gcagcacgag taatcgttgc 1860
 agtggccgag aatagaagcg gtttaactct tacttttgat aagccaaccg aggagaatgt 1920
 ggctaaggct cttaaaagca cggcgtctga ggccgtggta tgtcttgaac cgacatccga 1980
 agagggtgaac gtaaataaat tttctattgc tgagaaaggg agattgcctg tgtgtgcaga 2040
 aagtcatggt ttgacgaatg ctaacttaga gcaccaggag ttggagtccc tcaacgattt 2100
 ccataaggct tgcgtggata gtgtgattac aaagcaaatg gcatcggtt tctacactgg 2160
 ctactcaaa gttcaacaaa tgaagaacta tgtggacagt ttggcagctt cgttgtccgc 2220
 cactgtatca aatctatgca agtcactaaa ggatgaagtc gggatgatt ctgattccag 2280
 ggagaaagtt ggtgtttggg atgtcacttt gaaaaagtgg ctctctaaac ctgcggccaa 2340
 aggtcattca tggggagtgg tcttgatta caaggggaaa atgtttactg cacttctatc 2400
 ttatgaagga gatagaatgg tgactgagag cgactggagg aggggtggctg tatcatctga 2460
 tacaatggta tattctgata ttgcaaagct ccaaaatctg aggaaaacaa tgagagacgg 2520
 tgaacccac gaacctactg caaagatggt acttgtggat ggggtgcctg gttgtggaaa 2580
 gtacaaagga gattttgaaa gatttgatct tgatgaggat ttgatcttgg ttcttgaaa 2640
 acaagctgct gctatgatca gaagaagggc taattcatct ggactgataa gagccacaat 2700
 ggacaatgtg agaacggtag attcacttct aatgcatcca aaaccgcgat cacacaagag 2760
 gctttttatt gatgaagggt tgatgctgca caccggttgt gttaacttcc tgggtgcttat 2820
 ctctggttgc gacatcgcat acatttacgg agatacacag cagattcctt tcattaacag 2880
 agttcagaat tccccgtatc ccaaactatt tgagaagctg caagtggatg aagttgagat 2940
 gaggaggacc aactgagat gccagggtga tgtgaatttt ttcctacaat cgaagtacga 3000
 aggagcgggtg acaaccactt caactgtaca acgatcggtc tcatctgaga tgataggcgg 3060
 taagggagta ctaaacagtg tttccaaacc actaaaaggg aaaattgtaa ctttactca 3120
 ggctgataaa tttgagttag aggagaaggg ctataagaat gtgaacaccg ttcagatgat 3180
 ccaaggagaa acctttgaag atgtgtcgtt ggtcagattg acggcaactc cactgactct 3240
 gatttccaag tcttccccgc atgttctagt cgctctgact agacacacaa agagcttcaa 3300

aagacttaaa attcaggggtg gctgatacca aaatcagcag tgggtgttcg tccacttaaa 6240
 tataacgatt gtcatatctg gatccaacag ttaaaccatg tgatgggtgta tactgtggta 6300
 tggcgtaaaa catcggagag gttcgaatcc tcccctaacc gccggtagcg gccca 6355

<210> 6

<211> 2346

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequenc : nucleotide
 sequence of the tomato phytoene desaturase (pds)
 encoding cDNA

<400> 6

cttttactag ttatagcatt cggtatcttt ttctgggtaa ctgccaaacc accacaaatt 60
 acaagtttcc atttaactct tcaacttcaa cccaacccaaa tttattttcct taattgtgca 120
 gaaccactcc ctatatcttc taggtgcttt cattcggttcc gaggtaagaa aagatttttg 180
 tttctttgaa tgctttatgc cactcgttta acttctgagg tttgtggatc ttttaggcga 240
 cttttttttt ttttgtatgt aaaatttgtt tcataaatgc ttctcaacat aaatcttgac 300
 aaagagaagg aattttacca agtatttagg ttcagaaatg gataattttc ttactgtgaa 360
 atatccttat ggcaggtttt actgttattt ttcagtaaaa tgcctcaaatt tggacttggt 420
 tctgctgtta acttgagagt ccaaggtagt tcagcttatc tttggagctc gaggtcgtct 480
 tctttgggaa ctgaaagtcg agatgggtgc ttgcaaaggga attcgttatg ttttgctggt 540
 agcgaatcaa tgggtcataa gttaaagatt cgtactcccc atgccacgac cagaagattg 600
 gttaaggact tggggccttt aaaggctcgt tgcattgatt atccaagacc agagctggac 660
 aatacagtta actatttggg ggctgcattt ttatcatcaa cgttccgtgc ttctccgcgc 720
 ccaactaaac cattggagat tgttattgct ggtgcagggt tgggtgggtt gtctacagca 780
 aaatatttgg cagatgctgg tcacaaaccg atactgctgg aggcaaggga tgttctaggt 840
 ggaaaggtag ctgcatggaa agatgatgat ggagattggt acgagactgg tttgcatata 900
 ttctttgggg cttacccaaa tattcagaac ctgtttggag aattagggat taacgatcga 960
 ttgcaatgga aggaacattc aatgatattt gcaatgccaa gcaagccagg agaattcagc 1020
 cgctttgatt tctccgaagc tttaccgcgt cctttaaatg gaatttttag catcttaaag 1080
 aataacgaaa tgcttacatg gccagagaaa gtcaaatttg caattggact cttgccagca 1140
 atgcttgagg ggcaatctta tgttgaagct caagatggga taagtgttaa ggactggatg 1200
 agaaagcaag gtgtgccgga cagggtgaca gatgaggtgt tcattgctat gtcaaaggca 1260
 ctcaacttta taaaccctga cgaactttca atgcagtga ttttgatcgc attgaacagg 1320
 tttcttcagg agaaacatgg ttcaaaaatg gccttttttag atggtaatcc tcctgagaga 1380
 ctttgcatgc cgattgttga acacattgag tcaaaaagggtg gccaaagtcag actgaactca 1440
 cgaataaaaa agattgagct gaatgaggat ggaagtgtca agagttttat actgagtgaac 1500
 ggtagtgcaa tcgagggaga tgcttttgtg tttgccgctc cagtggatat tttcaagctt 1560
 ctattgcctg aagactggaa agagattcca tattttccaaa agttggagaa gttagtgcga 1620
 gtacctgtga taaatgtaca tatatggttt gacagaaaac tgaagaacac atatgatcat 1680
 ttgctcttca gcagaagctc actgctcagt gtgtatgctg acatgtctgt tacatgtaag 1740
 gaatattaca accccaatca gtctatgttg gaattgggtt ttgcacctgc agaagagtgg 1800
 atatctcgca gcgactcaga aattattgat gcaacgatga aggaactagc aacgcttttt 1860
 cctgatgaaa tttcagcaga tcaaagcaaa gcaaaaatat tgaagtacca tgttgtcaaa 1920
 actccgaggt ctgtttataa aactgtgccg ggttgtgaac cctgtcggcc tttacaaaga 1980

tccccaatag agggggtttta tttagccggt gactacacga aacagaaata cttggcttca 2040
atggaaggcg ctgtcttatac aggaaagctt tgtgctcaag ctattgtaca ggattatgag 2100
ttacttggtg gacgtagcca aaagaagttg tcggaagcaa gcgtagttta gctttgtggt 2160
tattatttag cttctgtaca ctaaatttat gatgcaagaa gcgttgtaga caacatatag 2220
aagaagagtg cgaggtgaag caagtaggag aaatgtagg aaagctccta tacaaaagga 2280
tggcattgtg aagattagca tctttttaat cccaagtta aatataaagc atattttatg 2340
gaattc 2346

<210> 7

<211> 7096

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: nucleotide
sequence of the tobacco nitrate reductase (nia-2)
encoding cDNA

<400> 7

tacatacaag ggcgcgaaata aacttttttt aaagtaaagt tatatgaact tgcaatgaaa 60
gaggacctta acttggttgt ctttggtgct ttctgcaaatt ttcaccttaa cagcccat 120
gagattgatt tagttagtta taacaattag ttaaagtgtt gtgtaatttg aagaaaatat 180
ttggacgtgc tcgctgaaaa cattatactc ctatataata gaaatacttt ctgaaaagtt 240
ggtcttggtc aaaaacgtat aagagagttg gtcttctcat aaatagtcac tagctttctg 300
attttttttc actttctata tcacgtaaatt aggtactcaa atttgatatt tacaccaaac 360
aaatgaaaat aggatatgtg tttttcatac gtatatttat ctatcgtact taatgatata 420
tacatataca tataacctta ctttttgatt actaaaaatt taattatatt taatttggtg 480
aaatatcaga tgccacaaaa catttaccta gccactgttt ttgactacta aaaatttaatt 540
tatgttttagc ttgggtaaat atcagatgtc actaaacatt ttacctagcc attcctccga 600
aaagaaattg agaaggaaat tagagttagt ggagccataa taatgtttta tgtgaccata 660
actcgggtgaa aaccacggca agaataagaa acagctgtta aggctaacca acagctgcat 720
atctttaagc catttgctat taccacaaca tcgcatcttc ctctgatccc gaccctacgg 780
gcgtaaaaag tgtaaatcgt tagaattgtt ttatttattt tatgatgtca ctatttttta 840
aaatcaaaat taaattgggg tgtcgatttt tttgggtcct gcttatgtat agtatggcgc 900
tatggaggca ctgagagagt ccgaaacgtt tctatataag gccacccac gcattcaca 960
acttcgttcc caaacagaa aagaaaatca aatctcggag agagagagag agaaatat 1020
tgagagagaa atacagaaaa tctctcttcc ttctttcctt tttttttcaa tccccattca 1080
tattcttttt ttagaataat ctatggcggc atctgtcgaa aacaggcagt tcagtcacct 1140
agaagccggt ttatcccggt ctttcaagcc ccggtctgat tccccggtt gtggctgcaa 1200
cttcccttcg cccaacagta ctaatttcca aaagaaacca aattccacca tttacctga 1260
ttactcgtcg agtgaagacg acgatgatga tgacgaaaaa aatgagtacc ttcaaatgat 1320
taaaaaaggg aattcagagt tagagccatc tgttcatgac actagggacg aaggtaccgc 1380
tgataattgg attgaacgca acttttccat gattcgtctc accggaaagc atccatttaa 1440
ctccgaacca ccgttgaacc ggctcatgca ccacggcttt atcacaccgg tcccacttca 1500
ttacgttcgt aacctgggac cggttcccaa gggcacgtgg gatgactgga ccgtggaagt 1560
cacgggacta gtgaagcgtc ctatgaaatt cacaatggac cagttgggtta acgaattccc 1620
ttgtagagaa ttgcccggtta cgcttggttg tgctggcaat cgaaggaaag aacagaacat 1680
ggttaaacaa accattgggt tcaactgggg cgccgctgcc gtttcaacaa cgatatggcg 1740

cggcggtacc	ctccgcgctt	tgctaaaaacg	gtgcgggtgt	tttagcaaga	ataaaggggc	1800
gcttaatgtt	tgcttcgaag	gagctgatgt	gttgcccgga	ggtggtggtt	caaagtatgg	1860
aaccagcatt	aagaaggaat	ttgcaatgga	tccagcacga	gatatcatcg	tagcctacat	1920
gcagaacgga	gaaaaatttg	cacccgacca	cgggtttcca	gtacgaatga	taattccagg	1980
attcattgga	ggaagaatgg	tgaaatggat	aaagaggatt	atagtcacca	ccaagaatc	2040
agacagctat	tatcatttca	aggacaatag	agttcttcct	ccccatgttg	atgctgaact	2100
tgcaaatacc	gaaggtacgt	accgtaacta	tttcaattta	ttactccatt	tgttccaatt	2160
tatgtgaacc	tatttccttt	ttggtccgtt	caaaaaagaa	tgaacccttt	ctaaatttgg	2220
taacaattta	gcttaaactt	acaacttcac	ccttaatgag	aaacttttat	aaccacacaa	2280
ataccctggg	gcccatttgg	acttgtttag	gtcgacaaat	tccaaaagtt	ttattttttt	2340
cttaaacttc	gtgctcagtc	aaacaggttc	acgtaaattg	aaacgggagag	agtatcattt	2400
ttattaaggg	gtataaatat	atttttaatta	gttgagactt	gcacatacaa	gtaaaatatt	2460
tcttagaata	caaaatcaac	tgaaagctta	cttctaatta	tatggttttg	aatttttcctt	2520
tcaatgaagt	aaataaaaag	gaaacaatta	tattcaacgc	atgtagggtat	atggtcctgt	2580
cattatctca	aatcaaattg	tttaaagaca	aaggactttg	gaaacataga	attgtcagct	2640
ttatagtatt	ggagtactat	attagttagc	tgtttgcac	tattcataat	tggtctatct	2700
gtgtgcagca	tggtggtaca	agccagagta	tatcatcaat	gagcttaata	ttaactctgt	2760
cattacgacg	ccgtgtcatg	aagaaatttt	gccaatatac	gcctggacga	ctcagcgacc	2820
ttacacgttg	aggggctatt	cttattcttg	ttagtatttt	tatatatttcc	gattttgctg	2880
agaatatcat	atttcttagt	tttgtcgata	catcgtatcc	tctaactctg	acgtttttact	2940
tcgtccttat	gcaccacttt	acgtccttac	tttctcagac	agttttattga	tgaaaactac	3000
ttactatttt	cgacccgata	gcctcagcgt	ccttaattaa	atgtgatgtt	ttgaaagaga	3060
tattctctcc	cgtctatttt	aattaatttt	tggtctgttt	tatacgtggg	aatctatttt	3120
taacattaat	taatatagaa	atgaaccata	ttaatattat	taatttcttc	attgaaaata	3180
caacaaatac	tcttcggctc	ttactacaat	gacaattttg	aagaaaaata	attaattcct	3240
tcctaataatc	tgaaaaatca	aataattgtg	accataaaaa	aagggtcaaaa	aattaattaa	3300
aatgaactgg	agagagtaaa	ttagaaaaata	taattatagc	actagtaatt	aaagttatta	3360
gatgtcttct	ttaaaaagcg	tgtgaaaact	ttaaagacga	aatataatat	gaatattatc	3420
taataacttag	aaagtgtcaa	taattggtag	acaattttaa	ctatatacta	gttaaaaagt	3480
ctgtcaatac	aactatttagt	attgggggatt	agagagaata	gtagtaaaat	ggagtaattg	3540
gacgcgatgag	cttgggcatg	ctgattgctg	tcagcttggt	tgctaattgtg	aaaaagaaaa	3600
tagtaagaaa	aggccaacat	ggttttgttt	attttattat	gtggtagtac	acaaaaacct	3660
ggggagcttt	cctagtctctg	aagagtcggg	ctttggtagc	acaaaattaa	tagtatagta	3720
taccaagtga	atattaaatt	caattgtcta	aagcacggaa	tctttttgac	tacttttagtt	3780
cctgcattctt	gggttgccctc	aacaacaccc	tttattgaat	tattatagta	atgttcaata	3840
taatatacaa	ttagaaaaca	ctctaagtgg	tcactttata	tggatctagt	caatactatt	3900
tcttctaaac	aacgtgccta	attacttccc	actttccagt	acatgaccac	cattaagttt	3960
aattttttgtc	aattccttgt	gcaattggcc	cttcaaatga	gcagaagtgt	tacgtaggaa	4020
aactaacttc	agctactatt	ataggagtaa	acctgttagg	aaaagatgct	cgaggaactg	4080
acaaaaacttg	tagaataatt	agccattgta	ttgattgaaa	tactgattgt	gaacgtgtaa	4140
caaacaggcg	gagggaaaaa	agtaacgcga	gtagaagtga	cgttggtagg	aggagaaaca	4200
tggaagtta	gcacactaga	tcaccagag	aagcccacca	aatatggcaa	gtactgggtg	4260
tggtgctttt	ggtcactcga	ggttgagggtg	ttagacttgc	tcagtgtctaa	agaaattgct	4320
gttcgagctt	gggatgagac	cctcaatact	caacccgaga	agcttattttg	gaacgtcatg	4380
gtacgttcac	ttcttctttt	acctttattt	cttttaactt	ctatatacta	gcggtgtaaa	4440
gttatttttac	accataagtt	aacttacaaa	aatatgtaac	tattttact	acgagtgatg	4500
agggcaagaa	ggggtttaag	tatttgacaa	taaatgtaaa	ccctgcaatt	ttgttcctaa	4560
tttttttatcc	tttcaactct	ttgtgattgc	ttcattatct	agattcacag	agcacatgtg	4620

ttcacatgcc	aaaacaaaaa	actacaaaac	aaaaaaacttt	tcactagctt	tagtctaaga	4680
ttcccccttt	tttttttggg	aggtgtgtgg	tcatactcc	atagatcaat	tccagccact	4740
gacgtaccaa	accctgaaaa	ttcctagtag	ttatagcgac	gtacaatcat	ttcatattat	4800
gtaagcagag	acgtgatcac	atgaactaga	tgtgaatacc	acttgcccag	tccaccaggt	4860
caattcatct	agatgtgtaa	atcttgacac	cagcactggg	tcacttttat	aacactagca	4920
tttaacaaca	tttcatcctt	gaacattact	tgggctaatt	aataagtatt	tttttttata	4980
tactctaaaa	attgtaatta	cataaatgaa	tttaacttat	acacgctgac	aatgttacta	5040
attccacttt	ttacggacgg	ttatctatag	aaatcattta	ggtgaaacaa	ttctcttaca	5100
ctatgatcag	tgtagtagta	taatggttat	tacattttct	aaatattgtg	ctatgttgca	5160
atgttcaggg	aatgatgaat	aattgctggg	tccgagtaaa	gatgaatgtg	tgcaagcctc	5220
acaagggaga	gattggaata	gtgtttgagc	atccgactca	acctggaaac	caatcagggtg	5280
gatggatggc	gaaggagaga	catttgagga	tatcagcaga	ggcacctcaa	acactaaaga	5340
agagtatctc	aactccattc	atgaacacag	cttccaagat	gtactccatg	tccgagggtca	5400
ggaaacacag	ctctgctgac	tctgcttgga	tcatagtcca	tgggtcatatc	tatgacgccca	5460
cgcgtttctt	gaaagatcac	cctgggtggga	ctgacagcat	tctcatcaat	gctggcactg	5520
attgcactga	ggaatttgat	gcaattcatt	ctgataaggc	taagaagctc	ttggaggatt	5580
tcaggattgg	tgaactcata	actactgggt	acacctctga	ctctcctggc	aactccgtgc	5640
acggatcttc	ttccttcagc	agctttctag	cacctattaa	ggaacttggt	ccagcgcaga	5700
ggagtgtggc	cctaattcca	agagagaaaa	tcccatgcaa	actcatcgac	aagcaatcca	5760
tctcccatga	tgtaggaaa	tttcgatttg	cattgccctc	tgaggatcaa	gtcttgggct	5820
tgctgttg	aaaacatatc	ttcctctgtg	ccgttattga	cgataagctc	tgcatgcgcg	5880
cttacacgcc	tactagcacg	atcgatgagg	tggggtagctt	cgagttgggt	gtcaagatat	5940
acttcaaagg	aattcacctt	aaattcccca	atggaggggca	aatgtcacag	tatcttgatt	6000
ctatgccgtt	agggtcattt	ctcgacgtga	aagggtccatt	aggtcacatt	gaataccaag	6060
gaaagggaaa	tttcttagtt	catggcaaac	agaagtttgc	caagaagttg	gccatgatag	6120
cagggtggaac	aggaataact	ccagtgtatc	aagtcatgca	ggcaattctg	aaagatccag	6180
aagatgacac	agaaatgtat	gtgggtgtatg	ctaacagaac	agaggatgat	attttactta	6240
aggaagagct	tgattcatgg	gctgagaaaa	ttccagagag	ggttaaagtt	tggtatgtgg	6300
ttcaggattc	tattaaagaa	ggatggaagt	acagcattgg	ttttattaca	gaagccattt	6360
tgagagaaca	tatccctgag	ccatctcaca	caacactggc	tttggttgt	ggaccacctc	6420
ctatgattca	atgtgctggt	aatccaaact	tggagaagat	gggctatgac	attaaggatt	6480
ccttattgggt	gttctaattt	taaaaacaaa	acaatatctg	caggaataaaa	tttttttttt	6540
ccccctatca	gttgtagata	ttgtatttgg	tttatcaccc	ccatgtacta	cgtagtgttt	6600
gtagttctta	cattttttatt	ttttagaatt	tttttaaacc	ttaggatata	aaggttttct	6660
cttccaacaa	agtgattcct	tagggaagaa	atgtactgta	ctgtactagt	atgtctaagc	6720
cgaaagttgt	aatgtttacc	atgacaaatt	gtattcaatt	cctcatggaa	tagtaacatt	6780
gtgttcatgt	gtcttctctg	aagcgatcct	caaaatatca	atgtatatat	atagtaattg	6840
caaaccattg	ttccttttcc	cgatgtagtt	aactactcct	tcttttagctt	ctagtctctg	6900
gtgaatatatt	ttttttctat	aactctttta	ttaatacggc	cttaaataag	agaaaagttt	6960
aaaccacgaa	tatcattatg	cagacgtata	ggtaattaat	ctactttttg	aaaaaaaaatc	7020
tattttcttt	atgtgggtcct	tcaaaaataat	attctagaac	cttttgtata	ttccctttta	7080
acttctattt	agtttt					7096

<210> 8

<220>

<223> Description of Artificial Sequence: nucleotide
sequence of the tobacco nitrite reductase (nir-1)
encoding cDNA

<400> 8

tttctattaa atttctggca ccttcattgc caaatccagc tagattttcc aagaatgctg 60
tcaagctcca cgcaactccg ccgtctgtgg cagcgccgcc agctgggtgt ccagagggtg 120
ctgctgagag gctagaaccc agagttgagg aaaaagatgg ttattggata ctcaaggagc 180
agtttagaaa aggcataaat cctcaagaaa aggtcaagat tgagaagcaa cctatgaagt 240
tgttcatgga aaatgggtatt gaagagcttg ctaagatacc cattgaagag atagatcagt 300
ccaagcttac taaggatgat attgatgtta ggcttaagtg gcttggcctc ttccatagga 360
gaaagaacca atatggggcg ttcatgatga gattgaagct tccaaatgga gtaacaacga 420
gtgcacagac tcgatacttg gcgagtgtga taaggaaata cgggaaagaa ggatgtgctg 480
atattacaac gaggcaaaat tggcagattc gtggagtgtg actgcctgat gtgcccgaga 540
tactaaaggg actagcagaa gttgggttga ccagtttgca gagtggcatg gacaatgtca 600
ggaatccagt aggaaatcct cttgctggaa ttgatccaga agaaatagta gacacagggc 660
cttactactaa tttgctctcc caatttatca ctggcaattc acgaggcaat cccgcagttt 720
ctaacttgcc aaggaagtgg aatccgtgcg tagtaggctc tcatgatctt tatgaacatc 780
cccatatcaa cgatctcgcg tacatgcctg ccacgaaaga tggacgattt ggattcaacc 840
tgcttgtggg tgggttcttc agcgcaaaaa gatgtgatga ggcaattcct cttgatgcat 900
gggttccagc tgatgatgtt gttccggttt gcaaagcaat actggaagct tttagagatc 960
ttggtttcag agggaacaga cagaaatgta gaatgatgtg gttaatcgat gaactgggtg 1020
tagaaggatt cagggcagag gtcgagaaga gaatgccaca gcaagagcta gagagagcat 1080
ctccagagga cttgggttcag aaacaatggg aaagaagaga ttatcttggt gtacatccac 1140
aaaaacaaga aggctacagc tttattgggtc ttcacattcc agtgggtcgt gttcaagcag 1200
acgatatgga tgagctagct cgtttagctg atgagtatgg ttcaggagag atccggctta 1260
ctgtggaaca aaacattatt attccaaca ttgagaactc aaagattgag gcactgctca 1320
aagagcctgt tctgagcaca ttttcacctg atccacctat tctcatgaaa ggtttagtgg 1380
cttgtagctg taaccagttt tgtggacaag ccataatcga gactaaagct cgttccctga 1440
tgataactga agaggttcaa cggcaagttt ctttgacacg gccagtgagg atgcactgga 1500
caggctgccc gaatacgtgt gcacaagttc aagttgcgga cattggattc atgggatgcc 1560
tgactagaga taagaatgga aagactgtgg aaggcgccga tgttttctta ggaggcagaa 1620
tagggagtga ttcacatttg ggagaagtat ataagaaggc tgttccttgt gatgatttgg 1680
taccacttgt tgtggactta ctagttaaca actttggtgc agttccacga gaaagagaag 1740
aaacagaaga ctaataaaat ttagaatagt tgggtatttt gctgtgttca taacatgtaa 1800
tgtatgataa atcaatgcaa acatttctac ctacgtgag 1839

<210> 9

<211> 1294

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA of the
beta-1,3-glucanase of *Nicotiana plumbagenifolia*

<400> 9

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial

Sequence:beta-glucuronidase encoding region

<400> 11

atggtccgtc ctgtagaaac cccaacccgt gaaatcaaaa aactcgacgg cctgtgggca 60
ttcagtctgg atcgcgaaaa ctgtggaatt gatcagcggt ggtgggaaag cgcgttataa 120
gaaagccggg caattgctgt gccaggcagt tttaacgata agttcgccga tgcagatatt 180
cgtaattatg cgggcaacgt ctggtatcag cgcgaagtct ttataccgaa aggttgggca 240
ggccagcgta tcgtgctgctg tttcgatgct gtcactcatt acggcaaagt gtgggtcaat 300
aatcaggaag tgatggagca tcagggcggc tatacgccat ttgaagccga tgtcacgccc 360
tatgttattg ccgggaaaag tgtacgtatc accgtttgtg tgaacaacga actgaactgg 420
cagactatcc cgccgggaat ggtgattacc gacgaaaacg gcaagaaaaa gcagtcttac 480
ttccatgatt tctttaacta tgccgggaatc catcgacgag taatgctcta caccacgccc 540
aacacctggg tggacgatat caccgtgggt acgcatgtcg cgcaagactg taaccacgcg 600
tctgttgact ggcagggtgt ggccaatggt gatgtcagcg ttgaactgct tgatgcggat 660
caacagggtg ttgcaactgg acaaggcact agcgggactt tgcaagtggg gaatccgcac 720
ctctggcaac cgggtgaagg ttatctctat gaactgtgct tcacagccaa aagccagaca 780
gagtgtgata tctaccgctc tcgcgtcgcc atccggtcag tggcagtga gggcgaacag 840
ttcctgatta accacaaacc gttctacttt actggctttg gtcgtcatga agatgcggac 900
ttacgtggca aaggattcga taacgtgctg atggtgcacg accacgcatt aatggactgg 960
attggggcca actcctaccg tacctcgcat tacccttacg ctgaagagat gctcgactgg 1020
gcagatgaac atggcatcgt ggtgattgat gaaactgctg ctgtcggctt taacctctct 1080
ttaggcattg gtttcgaagc gggcaacaag ccgaaagaac tgtacagcga agaggcagtc 1140
aacggggaaa ctcagcaagc gcacttacag gcgattaaag agctgatagc gcgtgacaaa 1200
aaccaccaa gcgtgggtgat gtggagtatt gccaacgaac cggatacccg tccgcaagtg 1260
cacgggaata tttcgccact ggcggaagca acgcgtaaac tcgacccgac gcgtccgcat 1320
acctgcgtca atgtaatgtt ctgcgacgct cacaccgata ccatcagcga tctctttgat 1380
gtgctgtgcc tgaaccgtta ttacggatgg tatgtccaaa gcggcgattt ggaaacggca 1440
gagaaggtagc tggaaaaaga acttctggcc tggcaggaga aactgcatca gccgattatc 1500
atcaccgaat acggcgtgga tacgttagcc gggctgcact caatgtacac cgacatgtgg 1560
agtgaagagt atcagtgtgc atggctggat atgtatcacc gcgtctttga tcgcgtcagc 1620
gccgtcgtcg gtgaacaggt atggaatttc gccgattttg cgacctcgca aggcatattg 1680
cgcgttggcg gtaacaagaa agggatcttc actcgcgacc gcaaaccgaa gtcggcggtc 1740
tttctgctgc aaaaacgctg gactggcatg aacttcggtg aaaaaccgca gcagggagggc 1800
aaacaatga 1809

<210> 12

<211> 411

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
part of the region of a TMV-U2 variant comprising

the origin of assembly

<400> 12

ccctcgccaa ttgaactcac tgaaaaagtt gttgatgagt tcgtagatga agtaccgatg 60
gctgtgaaac tcgaaagggt ccggaataca aaaaagagag tggtaggtaa taatgttaat 120
aataagaaaa taaataatag tggtaagaag ggtttgaaag ttgaggaaat tgaggataat 180
gtaagtgatg acgagtctat cgcgtcatcg agtacgtttt aatcaatatg ccttatacaa 240
tcaactctcc gagccaattt gtttacttaa gttccgctta tgcagatcct gtgcagctga 300
tcaatctgtg tacaaatgca ttaggtaacc agtttcaaac gcaacaagct aggacaacag 360
tccaacagca atttgcggtat gcctggaaac ctgtgcctag tatgacagtg a 411

<210> 13

<211> 198

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
STMV leader region

<400> 13

agtaaaactt accaatcaaa agacctaac aacaggactg tcgtgggtcat ttatgctgtt 60
gggggacata gggggaaaac atattgcctt cttctacaag aggccttcag tcgccataat 120
tacttggcgc ccaattttgg gtttcagttg ctgtttccag ctatggggag aggtaagggt 180
aaaccaaacc gtaaactg 198

<210> 14

<211> 455

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
STMV trailer region

<400> 14

gacaagtgcg cttgggttatt tcgtgttggt ttaactgaac ctgcacataa gccttttggg 60
tcgaagggtta aacgatccgc tcctcgcttg agcttgaggc ggcgatatct ttatgtcaac 120
agagacactt tgggtctatgg ttgtataaca atagatagac tcccgtttgc aagattaggg 180
ttaacagatc ttgccgttag tctggttagc gcgtaaccgg ccttgattta tggaatagat 240
ccattgtcca atggcctttgc caatggaacg ccgacgtggc tgtataatac gtcgttgaca 300
agtacgaaat cttgttagtg tttttccctc cacttaaatc gaagggtttt gttttggtct 360
tcccgaacgc atacgttagt gtgactaccg ttgttcgaaa caagtaaaac aggaaggggg 420
ttcgaatccc tccctaaccg cgggtaagcg gccca 455

<210> 15

<211> 1971

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
part of the genome of a TMV-U1 variant, comprising
MP and CP genes

<400> 15

ggaaacactg tgattatagc tgcattgttg gctcgtatgc ttccgatgga gaaaataatc 60
aaaggagcct tttgtggtga cgatagtctg ctgtacttcc caaagggttg tgagtttccg 120
gatgtgcaac actccgcgaa tcttatgttg aattttgaag caaaactgtt taaaaaacag 180
tatggatact tttgcggaag gtatgtaata catcacgaca gaggatgcat tgtgtattac 240
gatccccctaa agttgatctc gaaacttggg gctaaacaca tcaaggattg ggaacacttg 300
gaggagtcca gaagggtctc ttgtgatgtt gctgtttcgt tgaacaattg tgcgtattac 360
acacagttgg acgacgctgt atgggagggt cataagaccg cccctccagg ttcgtttgtt 420
tataaaagtc tgggtgaagta tttgtctgat aaagttcttt ttagaagttt gtttatagat 480
ggctctagtt gttaaaggaa aagtgaatat caatgagttt atcgacctga caaaaatgga 540
gaagatctta cgcgtgatgt ttaccctgtt aaagagtgtc atgtgttcca aagttgataa 600
aataatgggt catgagaatg agtcattgtc agaggtaaac cttctcaaag gagttaagct 660
tattgatagt ggatacgtct gtttagccgg tttggctcgt acgggcgagt ggaacttgcc 720
tgacaattgc agaggagggt tgagcgtgtg tctgggtggac aaaaggatgg aaagagccga 780
cgaggccact ctccgatctt actacacagc agctgcaaag aaaagatttc agttcaaggt 840
cgttcccaat tatgtataaa ccaccagga cgcgatgaaa aacgtctggc aagttttagt 900
caatattaga aatgtaaaga tgtcagcggg tttctgtccg ctttctctgg agtttgtgtc 960
ggtgtgtatc gtttatagaa ataataaaa attaggtttg agagagaaga tcacaagtgt 1020
gagagatgga gggcccatgg aacttacaga agaagttgtt gatgagttca tggaagatgt 1080
ccctatgtca atcaggcttg caaagtttcg atctcgaacc ggaaaaaaga gtgatgtccg 1140
taaagggaaa attagtagta gtgatcgggt agcgccgaac aagaactata gaaatgttaa 1200
ggattttgga ggaatgagtt ttaaaaagaa taatttaatc gatgatgatt cggagactac 1260
tgtcgccgaa tcggattcgt tttaaatatg tcttacagta tcactactcc atctcagttc 1320
gtgttcttgt cagcagcgtg ggccgaccca atagagttaa ttaatttatg tactaatgcc 1380
ttaggaaatc agtttcaaac acaacaagct cgaactgtcg ttcaaagaca attcagttag 1440
gtgtggaaac cttcaccaca agtgactgtt aggttccttg acagtgactt taagggtgtac 1500
aggtacaatg cggattatga cccgctagtc acagcactgt taggtgcatt tgacactaga 1560
aatagaataa tagaagttga aaatcaggcg aacccacaa ctgccgaaac gtttagatgct 1620
actcgtagag tagacgacgc aacggtggcc ataaggagcg ctataaataa tttagtagta 1680
gaattgatca gaggaaccgg atcttataat cggagctctt tcgagagctc ttctggtttg 1740
gtttggaaat ctggtcctgc aacttgaggt agtcaagatg cataataaat aacggattgt 1800
gtccgtaatc acacgtggtg cgtacgataa cgcatagtgt ttttccctcc acttaaatcg 1860
aagggttggt tcttggtatg cgcgggtcaa atgtatatgg ttcatataca tccgcaggca 1920
cgtaataaag cgaggggttc gaatcccccc gttacccccg gtagggggccc a 1971